



TASMANIAN SCALEFISH FISHERY ASSESSMENT 2013/14

Timothy Emery, Justin Bell, Jeremy Lyle and Klaas Hartmann

December 2015

TASMANIAN SCALEFISH FISHERY ASSESSMENT 2013/14

Timothy Emery, Justin Bell, Jeremy Lyle and Klaas Hartmann

December 2015



This assessment of the Tasmanian Scalefish Fishery is produced by the Institute for Marine and Antarctic Studies (IMAS) and uses input from the Scalefish Fishery Advisory Committee (SFAC). The SFAC met on the 26 November 2015 to consider the results of the 2013/14 Scalefish assessment. The SFAC participants were:

Stephen Bartels

Shane Bevis

Dave Blake

Jon Bryan

Todd Francis

Craig Garland

Ashley Kent

Jeremy Lyle

Tim Emery

Klaas Hartmann

Craig Midgely

Tim Mirabella

Grant Pullen

John Sansom

Frances Seaborn

Julian Harrington

The authors do not warrant that the information in this document is free from errors or omissions. The authors do not accept any form of liability, be it contractual, tortious, or otherwise, for the contents of this document or for any consequences arising from its use or any reliance placed upon it. The information, opinions and advice contained in this document may not relate, or be relevant, to a reader's particular circumstance. Opinions expressed by the authors are the individual opinions expressed by those persons and are not necessarily those of the Institute for Marine and Antarctic Studies (IMAS) or the University of Tasmania (UTas).

IMAS Fisheries and Aquaculture
Private Bag 49
Hobart TAS 7001
Australia

Email: Timothy.Emery@utas.edu.au

Ph: 03 6227 7284, +61 3 6227 7284 (international)

Fax: 03 6227 8035

© Institute for Marine and Antarctic Studies, University of Tasmania 2015

Copyright protects this publication. Except for purposes permitted by the Copyright Act, reproduction by whatever means is prohibited without the prior written permission of the Institute for Marine and Antarctic Studies.

Contents

Executive Summary	6
1. Fisheries Assessment.....	8
Management objectives and strategies	8
Major objectives	8
Primary strategies.....	8
The fishery.....	9
Data sources	9
Tasmanian Catch and effort logbooks.....	9
Commonwealth catch returns	9
Data analysis	10
Recreational fishery.....	13
Current assessment categories.....	14
Proposed assessment categories	15
Species' importance	15
Reporting levels definitions	15
Stock status definitions.....	17
Performance indicators and reference points definitions.....	18
2. General fishing trends.....	20
Commercial fishing licences.....	20
Commercial catch trends	20
General production	20
Estuarine production	24
Recreational fishery.....	26
Catch and effort.....	26
Recreational gillnet fishery.....	27
Uncertainties and implications for management	28
3. Gears in the commercial fishery	30
General effort trends.....	30
Automatic squid-jig.....	31
Beach seine	32
Drop-line	33
Dip-net.....	34
Danish seine.....	35
Fish trap.....	36

Graball net	37
Hand line.....	38
Small mesh net.....	39
Purse seine.....	40
Squid-jig.....	41
Spear	42
Trolling.....	43
Hand collection	44
4. Eastern Australian Salmon	45
5. Banded Morwong	52
6. Barracouta	65
7. Bastard Trumpeter	70
8. Blue Warehou	76
9. Flathead.....	83
10. Flounder	91
11. Gould’s Squid	97
12. Jack Mackerel	104
13. Jackass Morwong.....	111
14. Leatherjacket	118
15. Longsnout Boarfish.....	124
16. Yelloweye Mullet.....	129
17. Pike.....	134
18. School Whiting	139
19. Southern Calamari.....	145
20. Southern Garfish	153
21. Striped Trumpeter.....	162
22. Wrasse.....	174
References	181
Appendix 1- Common and scientific names of species from catch returns.....	189
Appendix 2- Data restriction and adjustments.....	190
Appendix 3- Annual Tasmanian scalefish production.....	193
Appendix 4- Annual catch, effort and number of vessels by fishing methods	195

Executive Summary

The Tasmanian scalefish fishery is a multi-species fishery operating in State fishing waters and encompassing a wide variety of capture methods. The Scalefish Management Plan (amended in 2009), provides the management framework for the fishery. A policy document includes the explicit identification of performance indicators and reference points that have two primary functions: (1) monitor performance of the fishery in relation to catch and effort, and (2) provide reference points against which the status of fish stocks can be assessed.

Fishery assessment

Since the early 1990's, annual commercial catches of the major species have generally been declining. Total scalefish production (excluding small pelagics) declined from over 2,000 t in the early 1990's to around 500 t in recent years due in part to changed targeting practices, the transfer of shark fishery to the Commonwealth and introduction of a State management plan in 1998.

There has been a continuous decline in the number of vessels participating in the scalefish fishery and in the number of scalefish fishing licences since 2000. Although effort performance indicators were not triggered for any fishing method, there are on-going concerns regarding the level of latent capacity within the fishery from licence holders who are currently either participating at low levels or not active (only 20-50% of licences are active depending on the type).

Catch and effort information for the recreational fishery are available periodically and have demonstrated that the recreational catch now represents a significant component of the total Tasmanian harvest for many species such as Flathead, Striped Trumpeter, Bastard Trumpeter, Blue Warehou, Flounder, Mullet, Barracouta, Jackass Morwong, Cod, Leatherjacket, and Silver Trevally.

Species assessment

The scalefish policy document is currently undergoing a review, including a review of performance indicators and reference points. For this assessment, a hierarchical approach based on the available information for individual species is applied in determining proposed reference points. Both existing and proposed reference points are, however, presented for each species.

The current assessment report should be viewed as transitional, both in terms of the development and implementation of reference points and the adoption of stock status reporting based on the recently implemented national stock status reporting framework.

Species/Species group	Status	Comment	Page
Australian Salmon <i>Arripis trutta</i>	SUSTAINABLE	Species has a long history of exploitation. Commercial landings are driven by market demand not abundance.	45
Banded Morwong <i>Cheilodactylus spectabilis</i>	TRANSITIONAL DEPLETING	Unusual combination of high longevity, fast growth and early maturity. The commercial fishery has been managed using Total Allowable Commercial Catch since 2008 and is closely monitored.	52
Barracouta <i>Thyrstites atun</i>	NOT ASSESSED	Large fishery for Barracouta in the 1960's but little market for the species nowadays.	65

Bastard Trumpeter <i>Latridopsis forsteri</i>	TRANSITIONAL DEPLETING	Catch rate and catches suggest record low population levels. The current minimum legal size limit is below the size of maturity.	70
Blue Warehou <i>Seriolella brama</i>	OVERFISHED	Commonwealth-managed species. Sporadically occurs in Tasmania. Despite a reduction in Total Allowable Catch (TAC) to 118 tonnes and introduction of a recovery plan for the species, the stocks have shown little signs of recovery.	76
Tiger Flathead <i>Platycephalus richardsoni</i>	SUSTAINABLE	Shared stock with Commonwealth fishery.	83
Sand Flathead <i>Platycephalus bassensis</i>	NOT ASSESSED	Sand Flathead is predominantly taken by recreational fishers. A recently introduced monitoring project through IMAS should allow an individual assessment to occur in 2014/15.	83
Flounder <i>Pleuronectidae</i> family	NOT ASSESSED	Greenback Flounder (<i>Rhombosolea tapirina</i>) constitute the majority of the commercial catches.	91
Gould's Squid <i>Nototodarus gouldi</i>	SUSTAINABLE	High inter-annual variability in abundance. Commonwealth fishery operating in Bass Strait. Dual-endorsed vessels fish in Tasmanian waters especially in years of peak abundance years.	97
Jack Mackerel <i>Trachurus declivis</i>	SUSTAINABLE	Shared stock with Commonwealth fishery, large biomass but low current catches.	104
Jackass Morwong <i>Nemadactylus macropterus</i>	SUSTAINABLE	Commonwealth-managed fishery.	111
Leatherjacket <i>Monacanthidae</i> family	UNDEFINED	Leatherjackets are a by-product species and are not actively targeted. There is little biological information available for most species.	118
Longsnout Boarfish <i>Pentaceropsis recurvirostris</i>	NOT ASSESSED	Little information is available for this species. Very low levels of catch. A trip limit applies	124
Mullet <i>Aldrichetta forsteri</i>	NOT ASSESSED	Recreational and commercial catches are similar and both minor.	129
Pike <i>Dinolestes lewini</i> (Longfin Pike) <i>Sphyaena novaehollandiae</i> (Shortfin Pike)	NOT ASSESSED	The main species targeted is the Shortfin Pike. Little information is available on either of the species.	134
School Whiting <i>Sillago flindersi</i>	SUSTAINABLE	Fishery currently based on one part-time operator.	139
Southern Calamari <i>Sepioteuthis australis</i>	UNDEFINED	Vulnerability of calamari to fishing pressure is unclear but probably high. There is increasing catch and effort in northern areas of the State and keen recreational interest in this species.	145
Southern Garfish <i>Hyporhamphus melanochir</i>	SUSTAINABLE	After a strong decline in catches from 2006/07 coupled with changes in population structure, which prompted management actions, the species is showing signs of recovery.	153

Striped Trumpeter <i>Latris lineata</i>	TRANSITIONAL RECOVERING	After a lack of recruitment for over a decade, Striped Trumpeter is showing signs of recovery in the last 3 years. However, the stock currently relies exclusively on recent recruits.	162
Wrasse <i>Notolabrus tetricus</i> (Blue-throated Wrasse) <i>Notolabrus fuciola</i> (Purple Wrasse)	NOT ASSESSED	Both species are targeted as part as a live fishery. A recent ban on the preferred bait type in fish traps has led to a decline in the participation in this fishery. Data is confounded due to mixed reporting of two species.	174

1. Fisheries Assessment

Management objectives and strategies

The Scalefish Fishery Management Plan [*Fisheries (Scalefish) Rules 1998*] was first introduced in 1998 (DPIF 1998) and was reviewed in 2001, 2004 and 2009. The management plan provides the regulatory framework for the fishery, which covers commercial and recreational components. While the plan contains the overarching legislation under which the fishery operates, the following objectives, strategies and performance indicators are contained in a policy document currently under review.

Major objectives

- To maintain fish stocks at sustainable levels by restricting the level of fishing effort directed at scalefish, including the amount and types of gear that can be used;
- To optimise yield and/or value per recruit;
- To mitigate any adverse interactions that result from competition between different fishing methods or sectors for access to shared fish stocks and/or fishing grounds;
- To maintain or provide reasonable access to fish stocks for recreational fishers;
- To minimise the environmental impact of scalefish fishing methods generally, and particularly in areas of special ecological significance;
- To reduce by-catch of juveniles and non-target species; and
- To implement effective and efficient management.

Primary strategies

- Limit total commercial fishing capacity by restricting the number of licences available to operate in the fishery;
- Define allowable fishing methods and amounts of gear that can be used in the scalefish fishery;
- Monitor the performance of the fishery over time, including identification and use of biological reference points (or limits) for key scalefish species;
- Protect fish nursery areas in recognised inshore and estuarine habitats by prohibiting or restricting fishing in these areas;
- Employ measures to reduce the catch and mortality of non-target or undersized fish; and
- Manage developing fisheries under permit conditions.

The fishery

The Tasmanian scalefish fishery is a multi-gear and multi-species fishery. The main gear types include gillnet, hooks and seine nets, harvesting a diverse range of scalefish, shark and cephalopod species. Other fishing gears in use include traps, Danish seine, dip nets and spears. A listing of common and scientific names of species reported in catches is presented in Appendix 1.

In many respects the scalefish fishery is dynamic, with fishers readily adapting and changing their operations in response to changes in fish availability and in response to market requirements and opportunities. As a consequence, only a small proportion of the fleet has specialised in a single activity or targeting a primary species (Ziegler 2012). For many operators, scalefish represent an adjunct to other activities, for instance rock lobster fishing.

This report covers the assessment of selected scalefish and cephalopod fisheries under Tasmanian jurisdiction, as well as selected species under Commonwealth jurisdiction. These were included due to the importance of catch levels in Tasmanian logbooks. Formal assessments of species under Commonwealth jurisdiction (e.g. Tiger Flathead, Blue Warehou, Jackass Morwong, Ocean Perch, Blue-eye Trevalla, Blue Grenadier, School and Gummy Shark) are undertaken by the Southern and Eastern Scalefish and Shark Fishery Assessment Group (SESSFAG, e.g. Morison et al. 2012) and are summarised in fishery status reports produced by the Australian Bureau of Agricultural and Resource Economics and Sciences (e.g. Woodhams et al. 2012).

Data sources

Commercial catch and effort data are collected through compulsory Tasmanian Commercial Catch, Effort and Disposal Returns, and Commonwealth non-trawl (GN01 and GN01A) and Southern Squid-jig Fishery (SSFJ) logbook returns. Unless noted otherwise, catch and effort data reported in this assessment relate to the commercial sector. Catch and effort information for the recreational sector are collected from surveys that are conducted periodically.

Tasmanian Commercial Catch, Effort and Disposal Returns

The catch and effort logbooks have been amended several times (1995, 1999, 2007, 2010 and 2013) in an effort to provide finer spatial scales and greater operational details. While the outer fishing blocks are still at the 30nm (1/2 degree) spatial resolution, the logbooks introduced in 2010 have redefined the scale of the coastal blocks (Fig. 1.1). In analysing catch and some data manipulation has been undertaken, details of which are provided in Appendix 2.

Commonwealth catch returns

Following the introduction of the Commonwealth non-trawl logbook (GN01 and subsequent versions) in late 1997, dual endorsed Tasmanian and Commonwealth (South East Non-Trawl and Southern Shark) operators generally commenced recording all of their catch and effort data, including fishing in State waters, in the Commonwealth logbooks. In addition, several dual endorsed squid operators reported some or all of their state waters fishing activity in the Southern Squid-jig Fishery (SSJF) logbook. As most of these operators did not explicitly indicate whether fishing occurred in State or Commonwealth waters, it has been necessary to incorporate all activity reported from coastal fishing blocks in the analyses. For details of data restrictions and manipulations involving Commonwealth logbook data see Appendix 2.

During 2001, dual endorsed fishers were instructed to report all fishing activities under State jurisdiction in the Tasmanian catch and effort logbooks. This should have removed the necessity to include subsequent Commonwealth catch and effort data into analyses, but it has become apparent that there was confusion amongst fishers about reporting requirements. For

example, catches of species such as Striped Trumpeter taken by Commonwealth operators were not routinely reported in the Tasmanian catch returns. Commonwealth logbook data since 2001 have been available for the current assessment and have been checked for possible double reporting (*i.e.* on both the Tasmanian and Commonwealth catch returns) and where this was not the case, the catch and effort database used in this assessment was updated.

Data analysis

For the purposes of this assessment, effort and catch rate analyses are restricted to commercial data provided for the period July 1995 to June 2014.

A fishing year from 1st July to 30th June in the following year has been adopted for annual reporting. This better reflects the seasonality of the fisheries for most species than reporting based on calendar year, with catch (and effort) generally concentrated between late spring and early autumn. In addition, it better encompasses the biological processes of recruitment and growth for most species. Unless stated otherwise, catches have been analysed state-wide and by region. Five broad assessment regions are used: south-east coast (SEC), east coast (EC), north-east coast including Flinders Island (NEC), north-west coast including King Island (NWC), and west coast (WC) (Fig. 1.1).




Figure 1.1 Map of Tasmania with the fishing blocks and the assessment regions. SEC is south-east coast, EC is east coast, NEC is north-east coast, NWC is north-west coast, and WC is west coast.

Two measures of effort have been examined in past assessments: (i) days fished (*i.e.* number of days on which a method/gear type was reported); and (ii) quantities of gear/time fished using the method. Since a diverse range of gear types are utilised in the fishery, appropriate measures of effort differ with gear type. Measures of effort by fishing method are presented in Table 1.1. However, because effort reporting has changed for some gears with the introduction of a new logbook during 2007, some gear related effort measures since 2007/08 are not directly comparable to those in the previous years. Additionally, the block reporting has changed in 2007 and some of the old block limits do not exactly match the new block subdivision. Consequently, confusion about the new reporting requirements may have biased some effort measures.

Table 1.1 Table of effort gear units by fishing method

Method	Effort gear units
Beach seine/purse seine	No. of shots
Graball/small mesh net	100 m net hours
Dropline	100 hook lifts
Handline	Line hours
Fish trap	No. trap or pot lifts
Squid-jig	Jig hours
Spear	Fisher hours
Dip-net	Dip-net hours

Catch returns for which *effort* information was incomplete or unrealistically high or low (either due to data entry error or misinterpretation of information requirements by fishers) were flagged and excluded when calculating effort levels based on gear units or catch rates based on catch per unit of gear. Only a small number of fishing records for 2013/14 needed to be excluded in this manner. All records were, however, included for reporting catch, days fished and catch per day.

In generating catch rate statistics, the geometric mean rather than the arithmetic mean of all valid individual daily catch records has been calculated, since catch rate data are typically log-normally distributed. The geometric mean is calculated as the n^{th} root of the product of the individual rates (y_i):

$$GM_{\bar{y}} = \sqrt[n]{\prod y_i}$$

This is equivalent to computing the arithmetic mean of the natural logarithm of each number, and then taking the exponent:

$$GM_{\bar{y}} = \exp \left[\frac{1}{n} (\sum \ln (y_n)) \right]$$

It should be noted that catch rates calculated in this manner may differ slightly from the more simplistic approach of dividing total catch by total effort or using the arithmetic mean. The geometric mean has the advantage of being less affected by the few observations that are skewed very high, as often happens with log-normally distributed data.

Recreational fishery

Information on recreational fisheries in Tasmania is relatively sparse in comparison to commercial data. Detailed analyses of the Tasmanian recreational fishery available are based on the 2000/01 National Survey (Lyle 2005), the 2007/08 state-wide fishing survey (Lyle et al. 2009) and the 2012/13 state-wide survey (Lyle et al. 2014b). Additional data are provided by a targeted study of offshore gamefish fishing (Tracey et al. 2013) and a targeted study of gillnet catch (Lyle and Tracey 2012) and fishing practices (Lyle et al. 2014a), along with recreational net licence numbers.

Current assessment categories

In the absence of rigorous quantitative stock assessments, the Scalefish Fishery Management Plan identifies a number of performance indicators that are used to define ranges between which the fishery, both in general and for particular species, is deemed to be performing acceptably (DPIF 1998). If the observed value of a performance indicator falls outside the acceptable range the reference point is said to have been exceeded implying that management action may be required. Analysis of fishery performance under this (initial) strategy is measured by reference to:

- variations in the total catch from year to year, or between seasons, regions and sectors;
- trends in effort;
- trends in catch rates;
- changes in biological characteristics, such as a changes in size or age structure; and
- other indicators of fish stock stress, for example disease outbreaks.

As part of this strategy, reference or trigger points, or acceptable ranges, have been defined as levels of, or rates of change, that are considered to be outside the normal variation of the stock(s) and the fishery. The trigger points provide a framework against which the performance of the fishery can be assessed and (if necessary) flag the need for management action. Currently, reference points for a species are exceeded when one or more of the following criteria are met:

Performance indicators	Reference points
Catch	<ul style="list-style-type: none"> • Total catch outside the 1990/91 to 1997/98 range • Total catch declines or increases by > 30% from previous year
Effort trend	<ul style="list-style-type: none"> • Effort (related to species) >10% higher than the maximum level from 1995/96 to 1997/98 • Effort (related to gear) >10% higher than the maximum level from 1995/96 to 1997/98
Catch rates trends	<ul style="list-style-type: none"> • Catch rate < 80% of lowest levels from 1995/96 to 1997/98
Change in biological characteristics	<ul style="list-style-type: none"> • Significant change in the size composition of commercial catches • Significant change in the abundance of a year class (or year classes) from size/age structure monitoring, with particular importance on pre-recruit year classes
Change in catch composition	<ul style="list-style-type: none"> • Change in the catch of non-commercial fish relative to 1995/96 to 1997/98 • Unacceptably high incidental mortality of non-commercial species or undersized commercial species
Stress	<ul style="list-style-type: none"> • Significant numbers of fish landed in a diseased or clearly unhealthy condition • Occurrence of a pollution event that may cause risks to fish stocks, the health of fish habitats or to human health • Any other indication of fish stock stress

Banded Morwong is the only scalefish species where management includes the use of a Total Allowable Catch (TAC), and therefore has its own set of performance indicators and reference points (see Banded Morwong section).

The fishery has also been assessed against an alternative set of performance indicators and reference points which account for recent developments in the fishery. These alternative assessment categories (presented below) are expected to be further refined and could replace the existing indicators over the next few years.

Proposed assessment categories

The proposed assessment categories take into account the species' importance and the quantity of data available to inform the appropriate reference points for each species, with the intention to assign a stock status to each species based on the presented criteria.

Species' importance

There are over 90 species reported in commercial catches. Harvested species in Tasmania have variable social and economic values, meaning that all species cannot be given the same priority for assessment. In assessing the importance of a species, a combination of factors was taken into account, which includes:

- Whether the species is a target, secondary target or by-product;
- The economic importance of the species;
- The annual landings for the species (i.e. annual catch >5 t for 50% of the time between 1995 and present);
- The number of operators targeting the species and;
- The "conservation" value of the species.

Species are consequently classified as "Key species" or "Minor species". The remaining species reported in commercial catches are considered of minor significance and not at threat from current fishing practices. As such, these species are not included in the assessment.

Reporting levels definitions

Each species in the assessment is associated with a level of reporting (Full, Medium or Minor), which stems from the available data for the species/species group. The aim is, in time, to have full reporting for all key species and medium reporting for all minor species. The attributes of the reporting categories are defined in Table 1.2. Table 1.3 summarises the species selected for the assessment, their importance and level of reporting.

Table 1.2. Summary of the attribute for the reporting categories

Attribute	Reporting level		
	Full	Medium	Minor
Time series estimate of biomass from dynamic models	•		
Time series estimate of total, natural and fishing mortality from dynamic models	•		
Quantitative risk analysis of future harvesting using dynamic models	•		
Time series of age and/or length composition data	•		
Estimates of total, natural and fishing mortality (from catch curves)	•		
Local (TAS) information for growth, mortality, selectivity and maturity	•	•	
Representative time-series of commercial catch	•	•	•
Single biological species or stock	•	•	•
Sporadic age and/or length composition data		•	
Non-local (non-TAS) information for growth, mortality, selectivity and maturity		•	•
Complex of related species		•	•



Table 1.3. Summary of importance and reporting level for all retained species

Species/Species group	Importance	Reporting level
Banded Morwong	Key	Full
Australian Salmon	Key	Medium
Bastard Trumpeter	Key	Medium
Blue Warehou	Key	Medium
Flathead	Key	Medium
Southern Calamari	Key	Medium
Southern Garfish	Key	Medium
Striped Trumpeter	Key	Medium
Wrasse	Key	Medium
Barracouta	Minor	Minor
Flounder	Minor	Minor
Gould's Squid	Minor	Minor
Jack Mackerel	Minor	Minor
Jackass Morwong	Minor	Minor
Leatherjacket	Minor	Minor
Longsnout Boarfish	Minor	Minor
Mullet	Minor	Minor
Pike	Minor	Minor
School Whiting	Minor	Minor

Stock status definitions

In order to assess the fisheries in a manner consistent with the national approach (and other jurisdictions) we have adopted the national stock status categories (Flood et al. 2012). These categories define the assessed state of the stock in terms of recruitment overfishing, which is often treated as a limit reference point. Recruitment overfishing occurs when the mature adult (spawning biomass) population is depleted to a level where it no longer has the reproductive capacity to replenish itself. Hence, recruitment overfished stocks have not collapsed but they do have reduced productivity. This is different to growth overfishing where fish are harvested at an average size that is smaller than the size that would produce the maximum yield per recruit. Fisheries are ideally also managed towards targets that maximise benefits from the harvesting, such as economic yield or provision of food. The scheme used here does not attempt to assess the fishery against any target outcomes.

Table 1.4 Stock status classification summary

Stock status	Description	Potential implications for management of the stock
SUSTAINABLE	Stock for which biomass (or biomass proxy) is at a level sufficient to ensure that, on average, future levels of recruitment are adequate (i.e. not recruitment overfished) and for which fishing pressure is adequately controlled to avoid the stock becoming recruitment overfished	Appropriate management is in place
TRANSITIONAL-RECOVERING 	Recovering stock—biomass is recruitment overfished, but management measures are in place to promote stock recovery, and recovery is occurring	Appropriate management is in place, and the stock biomass is recovering
TRANSITIONAL-DEPLETING 	Deteriorating stock—biomass is not yet recruitment overfished, but fishing pressure is too high and moving the stock in the direction of becoming recruitment overfished	Management is needed to reduce fishing pressure and ensure that the biomass does not deplete to an overfished state
OVERFISHED	Stock is recruitment overfished, and current management is not adequate to recover the stock; or adequate management measures have been put in place but have not yet resulted in measurable improvements	Management is needed to recover this stock; if adequate management measures are already in place, more time may be required for them to take effect
UNDEFINED	Not enough information exists to determine stock status	Data required to assess stock status are needed
NOT ASSESSED	No attempt has been made to assess stock status	Resources required to assess stock status

Performance indicators and reference points definitions

The determination of stock status is based on the consideration of model outputs (for species with full reporting) and the commercial catch and effort data, which are assessed by calculating fishery performance indicators and comparing them with reference points.

The proposed performance indicators are biomass and fishing mortality. For medium and minor reporting, proxies (commercial catch and CPUE) are used instead as there are insufficient data to calculate biomass or fishing mortality. These are compared to a reference period (1995/96¹ to 2006/07 unless stated otherwise) for each species. The reference points for more generic and full reporting are species-specific while the reference points for medium and minor reporting are applicable for all species. Reference points are illustrated in Fig. 1.2.

Table 1.5 Summary of the performance indicators and reference points for each reporting standard

Reporting	Performance indicators	Reference points			
Full	Fishing mortality	<ul style="list-style-type: none"> Maintain an appropriate spatial distribution of catch 			
	Biomass	<ul style="list-style-type: none"> High probability of staying above a certain level of mature biomass High probability of staying above a certain CPUE 			
Medium	Fishing mortality	<ul style="list-style-type: none"> Catch > 3rd highest catch value from the reference period Catch < 3rd lowest catch value from the reference period Catch variation from the previous year above the greatest inter-annual increase from the reference period Catch variation from the previous year above the greatest inter-annual decrease from the reference period Latest recreational catch estimate > recreational catch estimate from the reference period Proportion of recreational catch to total catch > previous proportion estimate 			
		Biomass	<ul style="list-style-type: none"> CPUE < 3rd lowest CPUE value from the reference period Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period 		
		Minor	Fishing mortality	<ul style="list-style-type: none"> Catch > 3rd highest catch value from the reference period Catch < 3rd lowest catch value from the reference period Latest recreational catch estimate > recreational catch estimate from the reference period Proportion of recreational catch to total catch > previous proportion estimate 	
				Biomass	<ul style="list-style-type: none"> Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period

¹ 1995/96 to 2006/07 was selected as the reference period, corresponding to the first twelve years since the introduction of daily catch and effort reporting in the Tasmanian General Fishing Returns.




Figure 1.2 Schematic illustration of the application of fishing mortality reference points for medium and minor reporting based on trends in catch. The shaded blue area represents the reference period. Note this figure is for illustrative purposes only.

As per the current assessment categories, other measures are also taken into consideration in the determination of stock status, such as changes in biological characteristics of the stock, indicators of stock stress, significant changes in the catch of non-commercial species or undersize commercial-species, and significant external factors related to fishing activity.

Reporting	Performance indicators	Reference points
All	Stock stress	<ul style="list-style-type: none"> • Significant change in the size/age composition of commercial catches • Significant numbers of unhealthy fish landed

As this is the third transitional report, stock status was only assessed for a few species (Australian Salmon, Bastard Trumpeter, Banded Morwong, Tiger Flathead, Gould’s Squid, Jack Mackerel, Leatherjacket, Southern Calamari, Striped Trumpeter and Blue Warehou) based on the current reference points. The proposed new reference points are also presented for each species for comparison.

2. General fishing trends

Commercial fishing licences

The number of scalefish licences A, B and C have slowly declined since 2000 to 296 total licences in 2013, of which 118 were active (Table 2.1). The decline was mainly due to a reduction in scalefish C fishing licences (which are non-transferable), while the numbers of scalefish A and B licences have remained stable. Scalefish C licences were also the licence type that was least often fished, with just 21% of C licences active in 2013 (down from 35% in 2000). The proportion of actively fished scalefish A and B licences dropped from around 70% up to 2007 to 54% for Scalefish A and 44% for Scalefish B licences in 2013.

In addition to the general scalefish licences, separate fishing licences allow the use of beach seine (a total of 50 licences in two categories A and B) and small mesh gillnet (10 licences). Fishers with a rock lobster licence (but without scalefish A or B licence) are also allowed to take scalefish with a limited amount of fishing gear.

Table 2.1 Number of active and total fishing licences (FL) by licence type (A, B or C) since 2000 (licence years run from March to February of the following year, hence the 2013 licencing year is the last complete year covered by this assessment).

No of active licences	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
FLA	50	44	51	48	46	38	43	47	37	33	32	35	28	35
FLB	109	104	111	110	109	101	105	105	93	88	84	76	80	67
FLC	79	62	63	52	47	34	36	33	23	16	17	17	13	16
Total	238	210	225	210	202	173	184	185	153	137	133	128	121	118

No of total licences	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
FLA	69	67	70	70	70	66	66	66	66	65	65	65	65	65
FLB	166	165	164	165	165	162	162	160	159	159	158	158	154	154
FLC	226	214	205	185	173	152	137	129	120	112	92	86	81	77
Total	461	446	439	420	408	380	365	355	345	336	315	309	300	296

Commercial catch trends

General production

Since the early 1990's, annual commercial catch of the major species has generally declined (Fig. 2.1). Overall, total scalefish production (excluding small pelagics and Gould's Squid) declined from over 2,000 t in the early 1990's to around 500 t in recent years (Appendix 3, Table A1).

The total scalefish catch increased by >100t between 2011/12 (499.8 t) and 2012/13 (610.7 t) but declined by just over 50% in 2013/14 (292.5 t). These large fluctuations are largely driven by the landings of Australian Salmon, which has the potential to mask changes in other species that are landed in much lesser quantities.

When assessing trends within the scalefish fishery it is important to recognise that some species occur seasonally in Tasmanian waters and that availability can differ markedly between years. Such variability may not necessarily reflect changes in stock status. Species in this category include Australian Salmon, Blue Warehou, Barracouta and Gould's Squid. By contrast, species such as Banded Morwong, Wrasse, Striped Trumpeter, Bastard Trumpeter, Southern Calamari and Octopus are 'resident' species, and variability in catches can reflect a combination of factors, including market preference, management intervention, stock status and intrinsic variability in life history.





Figure 2.1 Annual catches (t) for selected scalefish species since 1990/91. Tiger and Sand Flathead is combined.

 = Part Commonwealth-managed fisheries




Figure 2.1 Continued.




Figure 2.1 Continued.

Estuarine production

Estuarine production (a subset of the whole general fishery) totalled 62.7 t in 2013/14, which is an increase from the previous two years (Table 2.2). Catches came mainly from the southern estuaries of the Derwent River (ES1), Blackman Bay (ES17), Norfolk Bay (ES18), Frederick Henry Bay (ES19) and from the Tamar River (ES10). Whiting, Southern Calamari, Australian Salmon and Mullet were the main species. Port Davey (ES06) and Mersey River (ES08) are restricted to hand line fishing, which accounts for the very low production. Georges Bay (ES12) and Ansons Bay (ES11) has been closed to commercial scalefishing since 2004 and 2009 respectively.

Table 2.2 Total commercial catches (t) in estuaries around Tasmania (a) by fishing year and (b) by species in 2013/14

a) By fishing year	ES01	ES06	ES07	ES08	ES09	ES10	ES11	ES12	ES17	ES18	ES19	ES20	Total	ES	Description
95/96	30.8	0.8	4.4	0.2	0.7	11.4		0.4	2.9	26.4	16.1	3.2	97.3	ES01	Derwent River
96/97	37.7	0.3	2.6		0.6	16.6		0.9	6.4	12.3	7.0	1.8	86.2	ES06	Port Davey
97/98	59.5	0.2	1.4	<0.05	1.0	18.9		2.5	13.4	20.9	15.2	1.4	134.4	ES07	Macquarie Harbour
98/99	20.4		1.4		1.2	22.3		1.6	11.0	36.5	23.5	4.9	122.8	ES08	Mersey River
99/00	11.4		1.0		0.5	16.6	0.1	1.9	21.5	28.5	16.8	2.8	101.1	ES09	Port Sorrell
00/01	20.4		0.2		0.1	18.0	<0.05	1.2	16.0	28.0	28.4	1.9	114.2	ES10	Tamar River
01/02	14.0		2.2		0.2	81.9		1.2	9.2	64.6	32.6	2.0	207.9	ES11	Ansons Bay
02/03	30.4		8.1		0.2	29.8	0.5	0.8	14.6	35.4	23.4	1.6	144.8	ES12	Georges Bay
03/04	44.0		6.2		1.0	26.3			5.6	59.5	21.8	0.8	165.2	ES17	Blackman Bay
04/05	29.9		4.9		1.8	35.9			9.6	25.9	23.1	0.7	131.8	ES18	Norfolk Bay
05/06	4.1	0.1	23.2		0.9	34.2	1.3		8.4	14.4	19.6	0.8	107	ES19	Frederick Henry Bay
06/07	31.0	0.3	9.9		2.0	26.3	0.2		8.7	20.3	19.7	1.4	119.8	ES20	Pitt Water
07/08	38.8		3.2			17.9			13.0	26.9	12.3	0.9	113		
08/09	33.7		1.1		0.2	22.2			8.4	15.7	10.5	2.1	93.9		
09/10	26.9		0.7		0.5	17.5	<0.05		3.9	15.6	4.4	2.1	71.6		
10/11	27.8	0.1	0.4		0.6	11.1			6.3	5.4	13.7	1.7	67.2		
11/12	13.9		0.3			8.1			6.1	7.2	6.7	1.9	44.2		
12/13	12.3	0.1	0.1		<0.05	12.6	0.2		7.9	13.2	4.1	0.8	51.4		
13/14	32.3		1.1		0.3	9.9			2.6	6.7	8.7	1.1	62.7		
b) By species	ES01	ES06	ES07	ES08	ES09	ES10	ES11	ES12	ES17	ES18	ES19	ES20	Total		
Australian Salmon			1.0		0.1	1.4				<0.05	<0.05		2.5		
Blue Warehou											0.6		0.6		
Calamari	<0.05				0.2	1.0			1.3	1.0	2.4	<0.05	6.0		
Flathead	<0.05					<0.05			0.2	0.4	0.1		0.7		
Flounder	<0.05		<0.05			0.2			0.2	0.3	0.07	1.1	1.9		
Garfish						4.0				0.1	0.4		4.5		
Jack Mackerel						0.1							0.1		
Mullet						0.2							0.2		
Octopus						<0.05			<0.05	4.2	<0.05		4.3		
Other						<0.05							<0.05		
Sardines/Anchovy									<0.05	0.06			0.1		
Redbait													0.0		
Trevally	<0.05					<0.05							<0.05		
Whiting	32.2					<0.05					3.9		36.1		
Wrasse						<0.05				<0.05	0.1		0.1		
Total	32.3		1.0		0.3	7.0			1.7	6.8	7.6	1.1	57.8		

Recreational fishery

Catch and effort

Surveys of the recreational fishery conducted in 2000/01, 2007/08 and 2012/13 provide comprehensive snapshots of the Tasmanian recreational fishery (Henry and Lyle 2003, Lyle 2005, Lyle et al. 2009, Lyle and Tracey 2012, Lyle et al. 2014b). In addition there have been targeted surveys of recreational gillnetting in 1996-98 and 2010 (Lyle 1999, Lyle and Tracey 2012) and offshore boat fishing in 2011/12 (Tracey et al. 2013).

The most recent recreational fishing survey indicates that there has been a general decline in participation, in both absolute and relative terms (i.e. percentage of the population) (Lyle et al. 2014b). Nevertheless, recreational landings represent a significant component of the total harvest for many species, either as a proportion of the total harvest or in absolute quantities taken (Table 2.3). For instance, recreational catches exceeded commercial catches for Flathead, Black Bream², Mullet, Bastard Trumpeter, Jackass Morwong, Cod, Flounder, Barracouta and Silver Trevally during most survey years. By contrast, the commercial sector dominated the catches of Australian Salmon, Banded Morwong, Wrasse, Garfish, Whiting and Gould's Squid.

A particularly conspicuous change in the recreational fishery has been the more than threefold increase in the landings of Southern Calamari since 2000/01 such that the recreational harvest is now comparable to that for the commercial sector.

² It is illegal to retain Black Bream for commercial sale.

Table 2.3 Estimated recreational harvest (numbers and weight) for key scalefish species taken by Tasmanian residents, commercial landings (weight) and proportion of the total (recreational plus commercial) catch represented by the recreational harvest (refer Lyle et al. (2014b)). Note: the survey periods do not correspond with fishing years; 2000/01 represented the period May 2000 to Apr 2001, 2007/08 represented the period Dec 2007 to Nov 2008 and 2012/13 represented the period Nov 2012 to Oct 2013.* estimated from the 2011/12 offshore recreational fishing (Tracey et al. 2013); ** estimated from the 2010 recreational gillnetting survey (Lyle and Tracey 2012).

Species	2000/01				2007/08				2012/13			
	Rec harvest No.	Com. (t)	Rec. %	Rec. %	Rec harvest No.	Com. (t)	Rec. %	Rec. %	Rec harvest No.	Com. (t)	Rec. %	Rec. %
Flathead	1,236,675	322.0	63.4	83.5	1,066,293	293.0	73.2	80.0	924,932	235.9	39.9	85.5
Australian Salmon	300,456	105.0	485.0	17.8	110,312	48.1	299.8	13.8	144,712	63.7	270.8	19.0
Mullet	111,025	30.0	13.7	68.6	24,152	6.6	2.4	73.3	26,265	7.1	7.9	47.3
Flounder	50,582	15.2	10.5	59.1	32,436	10.1	7.8	56.3	23,238	7.2	2.1	77.4
Cod	65,115	30.6	4.0	88.4	14,263	8.2	2.5	76.7	10,464	6.1	2.2	73.5
Jackass Morwong	27,041	31.9	13.7	70.0	9,979	6.8	3.8	64.2	23,732	16.1	2.1	88.5
Garfish	15,669	1.9	81.4	2.3	14,568	2.0	51.0	3.7	15,260	2.0	50.6	3.8
Whiting	7,480	0.8	42.5	1.9	14,992	3.4	35.4	8.7	9,412	2.1	35.8	5.5
Black Bream	34,336	22.0	-	100.0	13,134	11.4	-	100.0	19,153	16.7	-	-
Barracouta	24,320	46.9	15.1	75.7	11,577	10.8	13.9	43.8	32,954	31.0	1.1	96.6
Wrasse	23,083	13.6	88.4	13.3	11,640	10.3	68.5	13.1	7,223	6.4	65.2	8.9
Blue Warehou	16,359	14.6	36.3	28.6	8,723	7.0	26.6	20.8	10,757	15.4	8.8	63.6
Jack Mackerel	15,770	3.2	8.6	26.8	5,216	1.0	225.7	0.4	30,907	5.2	0.2	96.3
Striped Trumpeter	13,450	29.6	49.6	37.4	7,274*	31.9*	19.8	61.7	3,476	15.2	10.5	59.1
Bastard Trumpeter	29,130	37.0	26.2	58.5	27,527**	27.3**	10.5	72.2	7,573	7.5	9.8	43.4
Leatherjackets	18,706	8.2	16.7	33.0	7,619	2.6	4.2	38.0	5,389	1.8	2.5	41.9
Silver Trevally	16,812	4.7	1.6	74.6	10,636	4.2	2.0	67.9	4,826	1.9	2.8	40.4
Southern Calamari	29,473	17.7	76.6	18.8	40,525	44.6	102.6	30.3	57,728	63.5	60.2	51.3
Gould's Squid	9,903	5.0	39.7	11.1	73,236	36.6	45.8	44.4	42,853	21.4	1054.7	2.0

Recreational gillnet fishery

The use of recreational nets in Tasmania has been subject to licensing since 1995, with fishers able to licence up to two graball nets prior to 2002, along with one mullet net and a beach seine. From November 2002 the number of graball nets that could be licensed was reduced to one per person. The number of recreational net licences issued rose rapidly from around 8 900 in 1995 to over 11 000 in 1999/00, licence numbers then stabilised at between 8 000–9 000 for several years. Licence numbers climbed again to around 10 000 in 2007/08 before trending downward to 8 606 by 2013/14 (Table 2.4). It is possible that the reduction in licence numbers since 2009/10 occurred in response to the introduction of maximum soak times for gillnets in 2009. Night netting, which was a common and popular practice amongst recreational fishers (Lyle 2000), was banned for recreational fishers (with the exception of Macquarie Harbour) in late 2004. While this appeared to have little discernible impact on licence numbers, a targeted survey of recreational gillnetting in 2010 revealed a concomitant reduction in overall gillnet effort (effort in 2010 was about 60% of the level in 1997 despite there being 40% more gillnet licence-holders, Lyle and Tracey 2012). Furthermore, only 73% of recreational licences were used during 2010 (Lyle and Tracey 2012).

The 2010 survey revealed that almost 65% of the gillnet catch (by number) was kept (Lyle and Tracey 2012). Bastard Trumpeter and Blue Warehou together represented 45% of the total

retained catch, Atlantic salmon contributed a further 10%, with Australian Salmon, Jackass Morwong, Mullet and Wrasse of secondary importance. Wrasse was the most significant by-catch. Recreational gillnet catches of Bastard Trumpeter, Mullet, Jackass Morwong, Leatherjacket and Cod were higher when compared to the commercial catches, while Blue Warehou catches were similar between the two sectors. Recreational gillnet catch rates have fallen from an average of >6 fish retained per net set in 1997 to just over 4 fish per set throughout the past decade. While variability in the abundance of target species has contributed to this trend (especially Blue Warehou), changes in fishing practices (no night netting, shorter average set durations, reduction in the length of mullet nets, larger minimum size limits for some species influencing release/discarding rates, etc.) have also been contributing factors.

Table 2.4 The number of recreational gillnet licences issued by licensing year since 1995/96. na = not applicable

Licence type	95/96	96/97	97/98	98/99	99/00	00/01	01/02	02/03	03/04	04/05	05/06
Graball 1	5615	6290	6685	6709	7477	7401	6960	7695	7313	7408	8054
Graball 2	2612	2678	2683	2426	2652	2515	1841	na	na	na	na
Mullet Net	656	684	738	739	879	845	608	754	753	754	816
Total	8883	9652	10106	9874	11008	10761	9409	8449	8066	8162	8870

Licence type	06/07	07/08	08/09	09/10	10/11	11/12	12/13	13/14
Graball 1	8677	9185	9172	8960	8162	8248	7995	7765
Graball 2	na	na	na	na	na	na	na	na
Mullet Net	877	995	1080	922	886	888	890	841
Total	9554	10180	10252	9882	9048	9136	8885	8606

Uncertainties and implications for management

While considerable attention has been directed at ensuring comparability of commercial data over time (refer Appendix 2), it is acknowledged that some recent administrative changes relating to the reporting of catches may have, nonetheless, exerted some influence on observed catch and effort trends.

Other uncertainties in this assessment relate to limitations in catch and effort data, mainly in terms of the level of detail provided and the lack of independent verification. Since the Commercial Catch, Effort & Disposal logbook (formerly the General Fishing Return) was designed to accommodate a diverse range of fishing activities, compromises have been necessary, with data collection on a daily rather than operational (set or shot) basis.

It has also become apparent that some fishers have experienced problems in correctly interpreting or complying with reporting requirements, especially in terms of how effort information is reported. The introduction of new logbooks during the 2007/08 season has helped to clarify reporting, but there continues to be an on-going need to educate fishers in this regard. Further, the lack of catch verification remains a major issue in relation to data quality. Anecdotal reports suggest that some catch and effort data may be unreliable, particularly prior to the implementation of the management plan in 1998.

Catch and effort are influenced by a combination of factors which include fishers matching their fishing operations to changing market requirements and/or resource availability, as well as responses to changing management arrangements. The latter adds further uncertainty regarding the underlying causes of any observed trends in catch and effort. There is, therefore, a need to take account of industry perceptions and information when interpreting fishery dependent information.

Limited information about the recreational fishery remains a major uncertainty and is especially significant in the scalefish assessment given the scale of recreational catches relative to commercial catches for some species. While the 2000/01, 2007/08 and 2012/13 surveys

provide important information about this sector, there is a need for an on-going monitoring program for key species in the recreational fishery. Without such information attempts to assess the status of those species with significant recreational catches may be compromised.

Fish mortality due to predation and fishery interactions with Australian and New Zealand fur seals is largely unknown but represents another source of uncertainty. Seals can cause substantial mortality to some of the fish species assessed in this report as well as causing gear damage and influencing fishers' behaviour, factors that impact on catches and catch rates. This tends to be caused predominantly by individual 'rogue' seals which learn to target particular fisheries or fishing methods (e.g. the Banded Morwong gillnet fishery), while the typical diet of seals includes mainly pelagic fish species (Goldworthy et al. 2003).

3. Gears in the commercial fishery

There are 14 main fishing methods used in the commercial Tasmanian scalefish fishery. Catch and effort by gear types are presented in Table A2 in Appendix 4. Since a variety of gear types are represented, it has been necessary to express effort in units appropriate to each specific fishing method (Table 1.1). Effort has also been expressed in terms of number of days fished using the specified gear type, irrespective of the amount of gear utilised each day. Although days fished represents a less sensitive measure of effort, it has become apparent that some fishers have misinterpreted reporting requirements for effort. Attempts have been made to reduce this problem by updating the logbook; however misreporting effort continues to be prevalent and due to logbook changes is inconsistent through time. Days fished overcomes any uncertainty about the reporting of effort units and provides a consistency providing there has been no major changes to daily fishing practices throughout the time series.

For the purpose of analysis, shark net and bottom longline catch and effort have been excluded since these methods relate specifically to the school and gummy shark fishery, which is managed by the Commonwealth.

General effort trends

Following the introduction of the new management arrangements in November 1998, effort based on beach seine, purse seine, gillnets and hand line all fell, whereas effort based on drop-line, squid-jig and dip-net increased sharply. While a range of factors, including availability of target species and market developments, have had an influence, there is little doubt that management changes have had a direct impact on effort levels. Specifically, there was a decline in the effort of fishing methods for which gear allocations, or access, became more regulated (beach seine, purse seine and gillnets), whereas there was an initial shift to or increase in effort for less regulated methods (hooks, jigs and dip-nets; i.e. gear that is equally available to all licence-holders).

Since the early 2000's effort levels for most fishing methods declined, exceptions being handline, which has remained relatively stable, and automatic squid-jig which has peaked sporadically with the periodic occurrence of Gould's Squid in Tasmanian waters. For example, catches were at a record high level during 2012/13 whereas no fishing was reported for the automatic jig fishery in Tasmanian waters during 2013/14.

In the other gear sectors, effort levels during 2013/14 were generally similar to, or lower than, those during 2012/13 (Figures 3.1–3.27; more detailed data available in Table A2, Appendix 4). There is, however, potential for future increases in some gear types due to the levels of latent effort from licence holders who are currently inactive in the fishery or participating at low levels. The 2004 management plan review in particular attempted to address this issue through several strategies including non-transferability of C-class licences.

The following section presents an overview of the catch composition, as well as the overall catch, effort, and number of vessels for each fishing method.

Automatic squid-jig

Automatic squid-jig users targets exclusively Gould's Squid, and have practically no by-catch.




Figure 3.1 Overall catch, number of vessels using the gear, and effort (in gear unit and days fished) for automatic squid-jig.

Beach seine




Figure 3.2 Beach seine catch composition for 2013/14.




Figure 3.3 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for beach seine.

Drop-line




Figure 3.4 Drop-line catch composition for 2013/14.




Figure 3.5 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for drop-line. For the purpose of analysis, drop-line catch and effort up to 1998 was restricted to records that indicated a fishing depth of less than 200 m, effectively excluding effort targeting Blue-eye Trevalla but encompassing the target fishery for Striped Trumpeter (less than 1% of the Striped Trumpeter catch has been reported from depths greater than 200 m; since 1998 fishing for Blue-eye has been covered in Commonwealth catch returns).

Dip-net




Figure 3.6 Dip-net catch composition for 2013/14.




Figure 3.7 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for dip-net.

Danish seine




Figure 3.8 Danish seine catch composition for 2013/14.




Figure 3.9 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for Danish seine.

Fish trap




Figure 3.10 Fish trap catch composition for 2013/14.




Figure 3.11 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for fish trap.

Graball net

Graball nets in this analysis include both the traditional graball net (110 mm mesh size) and the Banded Morwong net (140 mm mesh size).




Figure 3.12 Graball net catch composition for 2013/14.




Figure 3.13 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for graball net.

Hand line




Figure 3.14 Hand line catch composition for 2013/14.




Figure 3.15 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for hand line.

Small mesh net




Figure 3.16 Small mesh net catch composition for 2013/14.




Figure 3.17 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for small mesh net. Note: the low effort in 2010/11 is due to a significant amount of incorrect records for that fishing year.

Purse seine




Figure 3.18 Purse seine catch composition for 2013/14.




Figure 3.19 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for purse seine.

Squid-jig




Figure 3.20 Squid-jig catch composition for 2013/14.




Figure 3.21 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for squid-jig.

Spear




Figure 3.22 Spear catch composition for 2013/14.




Figure 3.23 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for spear.

Trolling




Figure 3.24 Trolling catch composition for 2013/14.




Figure 3.25 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for trolling.

Hand collection




Figure 3.26 Hand collection catch composition for 2013/14.





Figure 3.27 Overall catch (t), number of vessels using the gear, and effort (in gear unit and days fished) for hand collection.

4. Eastern Australian Salmon

Arripis trutta

STOCK STATUS	SUSTAINABLE
Species has a long history of exploitation. Commercial landings are driven by market demand.	
IMPORTANCE	Key
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends



Arripis trutta
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source												
Habitat	<ul style="list-style-type: none"> Open water. Down to 30 m depth. 	Edgar (2008)												
Distribution	<ul style="list-style-type: none"> From Victoria to Queensland (Brisbane) and around Tasmania. 	Edgar (2008)												
Diet	<ul style="list-style-type: none"> Fish (pelagic predominantly) 	Stewart et al. (2011)												
Movement and stock structure	<ul style="list-style-type: none"> One single well-mixed stock along south-east Australia Travel great distances between the different States Increased population mixing occurs with both increasing age and decreasing latitude 	Stewart et al. (2011)												
Natural mortality	<ul style="list-style-type: none"> M between 0.35 and 0.50. 	Stewart et al. (2011)												
Maximum age	<ul style="list-style-type: none"> Maximum sampled is 12 years but potentially up to 26 years. 	Stewart et al. (2011)												
Growth	<ul style="list-style-type: none"> Maximum length: 89.0 cm. Maximum weight: 9.4 kg Growth described by von Bertalanffy growth function $L = L_{\infty}(1 - e^{-k(t-t_0)})$ where L is the fork length (cm), t is the age (years), L_{∞} is the average maximum length for the species, k is a constant and t_0 is the (theoretical) age where length equals zero. Parameters estimates are: <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Sex</th> <th>L_{∞}</th> <th>k</th> <th>$t_{0\infty}$</th> </tr> </thead> <tbody> <tr> <td>Females</td> <td>65.0</td> <td>0.24</td> <td>-0.15</td> </tr> <tr> <td>Males</td> <td>61.3</td> <td>0.27</td> <td>-0.13</td> </tr> </tbody> </table> 	Sex	L_{∞}	k	$t_{0\infty}$	Females	65.0	0.24	-0.15	Males	61.3	0.27	-0.13	Edgar (2008) Frimodt (1995) Stewart et al. (2011)
Sex	L_{∞}	k	$t_{0\infty}$											
Females	65.0	0.24	-0.15											
Males	61.3	0.27	-0.13											
Maturity	<ul style="list-style-type: none"> Size-at-50% maturity: 42.13 cm Age at 50% maturity: 2.19 years 	Stewart et al. (2011)												
Spawning	<ul style="list-style-type: none"> From October to March, off New South Wales. The relationship between batch fecundity and fork length 	Stewart et al. (2011)												

	<p>is exponential with $F = 14581e^{0.0659L}$, where F is the fecundity (in number of eggs) and L is the fork length (cm).</p> <ul style="list-style-type: none"> The relationship between batch fecundity and age is exponential with $F = 96604e^{0.227A}$, where F is the fecundity (in number of eggs) and A is the age (in years). 	
Early life history	<ul style="list-style-type: none"> Eggs, larvae and juveniles drift and migrate from spawning grounds to Tasmania and Victorian waters during autumn and winter. Juveniles (4–6 cm fork length) appear in shallow Tasmanian waters between January and September. 	Kailola et al. (1993)
Gillnet post release survival	<ul style="list-style-type: none"> Low: 20 – 62% depending on soak duration 	Lyle et al. (2014a)

Background

There are two species of Australian Salmon cohabiting in Tasmanian waters: *Arripis trutta* (Eastern Australian Salmon) and *Arripis truttaceus* (Western Australian Salmon). The Eastern Australian Salmon constitute 94% of Tasmanian commercial catches.

Australian Salmon have a long history of exploitation in Tasmania, with large-scale commercial fishing occurring at least since 1958 (Stewart et al. 2011). There are two distinct sectors in the commercial fishery: a small number of large vessels specifically equipped to capture and store large quantities of Australian Salmon, and a large number of small vessels which target the species on an opportunistic basis or take them as by-product. One company operating up to three vessels accounts for around 85% of landings for the species.

Beach seining accounts for the majority of the catch. Large-scale beach seine operations involve deploying a net around a school of Australian Salmon using a small boat and then hauling the net into the shallows, forcing the fish to the cod-end which are then transferred to the mother ship. Spotter planes are typically used to locate the schools. Purse seine and gillnetting are often used by smaller operators. A total commercial catch trigger of 435 t (120% of the average annual catch between 1996/97 and 2006/07) applies to the Tasmanian fishery.

Most commercially caught Australian Salmon are frozen whole and sold as rock lobster bait with production levels linked to a large extent to the demand for bait. Some Australian Salmon are sold fresh for human consumption.

Australian Salmon is the second most commonly captured fish species by recreational fishers (Lyle 2005, Lyle et al. 2009, Lyle et al. 2014b), who mainly use line fishing methods to target the species.

FISHING METHODS	Mainly beach seine, also purse seine and gillnet. Line for recreational.
MANAGEMENT METHODS	<p>Input control:</p> <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence, Beach seine licence) Species licence (Australian Salmon licence) – 8 issued, 1 commonly used. Spatial and temporal area closures for Australian Salmon licence. <p>Output control:</p> <ul style="list-style-type: none"> Possession limit of 500kg on vessel at any one time for Scalefish fishing licence unless otherwise endorsed. Possession limit of 15 individuals for recreational

MAIN MARKET	<ul style="list-style-type: none"> • Minimum size (200 mm TL) • Total commercial catch limit of 435 t
	Local and interstate

Current assessment

Catch, effort and CPUE

Following a large catch in 2012/13 (331.3 t) landings were at a historic low of 38 t in 2013/14 (Fig. 4.1). The low catch was due to a dramatic decline in the landings by beach seine fishers that historically landed the majority of the catch. The majority of catch was caught on the north-east, north-west and south-east coasts, similar to previous years (Fig.4.1A and 4.2). Effort was low and similar to the level from the previous year. Catch rates by gear and days fished were the highest recorded since 1995/96 (Fig. 4.1C). It should be noted however that catch rate is influenced by the skewed nature of the data (i.e. the majority of catches are small but the total catch is influenced by only a small number of extremely large catches). In addition, catch rate is not a particularly sensitive indicator of fish stock condition for schooling species such as Australian Salmon, especially if search time is not taken into account.

Australian Salmon





Figure 4.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. BS= beach seine, NEC= north-east coast, NWC= north-west coast.

A) Catch



B) Effort




Figure 4.2 (A) Australian Salmon catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Total commercial catch >435 t	No	
	• Catch > higher catch from the 1990/91 to 1997/98 range (878.8 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (287.3 t)	Yes	249.6 t (86.9%)
	• Catch increases by > 30% from previous year (>430.7 t)	No	
	• Catch decreases by > 30% from previous year (<231.9 t)	Yes	194.2 t (83.7%)
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>835 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0195 t/days fished)	No	

Given that catch rates are not a significant indicator of stock status, the biomass performance indicators (based on CPUE and CPUE trends) were not calculated for Australian Salmon.

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Total commercial catch >435 t	No	
	• Catch > 3 rd highest catch value from the reference period (462.1 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (254.2 t)	Yes	216.5 (85.2%)
	• Catch variation from the previous year above the greatest inter-annual increase from the reference period (188.7 t)	No	
	• Catch variation from the previous year above the greatest inter-annual decrease from the reference period (240.1 t)	Yes	53.6 t (22.3%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period (105.2 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (13.8% in 2007/08)	Yes	Latest estimate (2012/13) 19.0%
Stock stress	• Significant change in the size/age composition	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

SUSTAINABLE

Although several catch and catch rate reference points were breached as a result of the low landings in 2013/14, annual production of Australian Salmon is strongly linked to the fishing practices of a single operator that is responsible for the overwhelming majority of landings. During the 2013/14 season, this operator was heavily involved in other fisheries, therefore Australian Salmon landings were greatly reduced. Most other beach seine fishers rarely, if at all,

target Australian Salmon. Due to the reduced catch, the Rock Lobster industry predominantly relied on Australian Salmon imported from New Zealand for bait. Due to these factors, the reduced catch is not representative of abundance.

Eastern Australian Salmon represent a single well-mixed stock along south-east Australia and the Tasmanian fishery catches mostly sub-adults. There has been little change in size and age composition in Tasmania or other states (NSW and VIC) suggesting the species is not overexploited. The eastern Australian biological stock is sustainably fished as a whole and the current level of commercial and recreational fishing pressure in Tasmania is well below historical levels and unlikely to cause the biological stock to become recruitment overfished.

5. Banded Morwong

Cheilodactylus spectabilis

STOCK STATUS	TRANSITIONAL DEPLETING
Unusual combination of high longevity, fast growth and early maturity. The commercial fishery has been managed using a Total Allowable Catch (TAC) since October 2008 and is closely monitored.	
IMPORTANCE	Key
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends; changes in size/age composition



Species biology

Parameters	Estimates	Source															
Habitat	<ul style="list-style-type: none"> Rocky reefs down to 50m depth. Females and juveniles inhabit the shallow section of the reef while males dominate in the deeper section of the reef. 	McCormick (1989a, b)															
Distribution	<ul style="list-style-type: none"> South Sydney (New South Wales) to eastern Victoria, Tasmania, New Zealand 	Gomon et al. (2008)															
Diet	<ul style="list-style-type: none"> Invertebrates, algae, crabs. 	McCormick (1998)															
Movement and stock structure	<ul style="list-style-type: none"> Limited movement of juveniles and adults, generally restricted to within 5km of the release site No information on the stock structure 	Murphy & Lyle (1999) Ziegler et al. (2006) Buxton et al. (2010)															
Natural mortality	<ul style="list-style-type: none"> Low. Estimated at $M=0.05$ 	Murphy & Lyle (1999)															
Maximum age	<ul style="list-style-type: none"> Females: 93 years Males: 96 years 	Ewing et al. (2007)															
Growth	<ul style="list-style-type: none"> Males grow to larger sizes than females Growth described by a Schnute-Richards growth function $L = L_{\infty}(1 + \alpha^{-at^c})^{-\frac{1}{b}}$ <p>where L is the length (mm), t is the age (years), L_{∞} is the average maximum length for the species and α, a, b and c are (year-specific) constants. The most recent parameters estimates (2007) are :</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>ex</th> <th>L_{∞}</th> <th>b</th> <th>c</th> <th>α</th> </tr> </thead> <tbody> <tr> <td>Female</td> <td>442</td> <td>18.8</td> <td>$3.3e^{-t}$</td> <td>0.05</td> </tr> <tr> <td>Males</td> <td>516</td> <td>2.3</td> <td>0.0088</td> <td>0.33</td> </tr> </tbody> </table> <ul style="list-style-type: none"> Length-weight relationship for 2007 was estimated at $W = 3.563E^{-5}L^{2.875}$ for females and $W = 3.729E^{-5}L^{2.852}$ for males, where W is weight (g) and L is the fork length (cm). 	ex	L_{∞}	b	c	α	Female	442	18.8	$3.3e^{-t}$	0.05	Males	516	2.3	0.0088	0.33	Schnute & Richards (1990) Ziegler et al.(2007a)
ex	L_{∞}	b	c	α													
Female	442	18.8	$3.3e^{-t}$	0.05													
Males	516	2.3	0.0088	0.33													

Maturity	<ul style="list-style-type: none"> Size-at-50% maturity estimated at 320 mm for females (~2.5 years of age). 	Ziegler et al.(2007a)
Spawning	<ul style="list-style-type: none"> Spawning occurs between mid-February to late May. Species is a serial spawner. 	Murphy & Lyle (1999)
Early life history	<ul style="list-style-type: none"> Eggs and larvae are concentrated on the surface. Banded Morwong has a pelagic stage distributed offshore, as suggested by the large amounts of larvae caught off the shelf break of eastern Tasmania. Juveniles appear in shallow water on rocky reefs and tide-pools between September and December, after a pelagic phase of 4-6 months. 	B. Bruce, Pers. Com. Wolf (1998)
Gillnet post release survival	<ul style="list-style-type: none"> High: 97% irrespective of gillnet soak duration 	Lyle et al. (2014a)

Background

The 'live fish' fishery for Banded Morwong began in the early 1990s. All holders of a fishing licence (vessel) were able to take this species and, as a result, there was a dramatic increase in effort directed at the species, with catches peaking at 145 t in 1993/94. Since then, catches have stabilised around 35–50 t. A quota management system with a Total Allowable Catch (TAC) was introduced in October 2008 (see Fig. 5.1 for areas). The TAC is currently undergoing a staged reduction as follows:

Quota year	TAC (in t)	TAC (in no. fish)	No. of Fish/Quota Unit
2012/13	38.8	29,825	25
2013/14	37.2	28,632	24
2014/15	35.7	27,439	23

Banded Morwong are targeted almost exclusively for the live fish market with large mesh gillnets, primarily 130–140 mm stretched mesh. The fishery is centred mainly along the east coast of Tasmania, between St. Helens in the north and the Tasman Peninsula in the south, with the largest catches traditionally coming from around Bicheno (Fig. 5.5). Smaller catches have been taken along the south coast and around Flinders Island. Fishing operations are conducted over inshore reefs, with gear set primarily in the 5–20 m depth range. Banded Morwong is not an important recreational species with recreational landings estimated between 1 and 2 t per year and being predominantly captured by gillnet. As such, when captured at any time other than by Banded Morwong fishers they constitute bycatch. They do, however, have exceedingly high post-release survival under current maximum permitted gillnet soak durations if carefully removed from gillnets (Lyle et al. 2014a).

FISHING METHODS	Mainly graball net
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence) Species licence (Banded Morwong licence) Temporal closure (March-April) Output control: <ul style="list-style-type: none"> Possession limit of 2 individuals for recreational fishers Minimum and maximum size (360-460 mm TL) TAC of 27,439 fish (35.7 t) for the 2014/15 licensing year
MAIN MARKET	Interstate




Figure 5.1 Designated TAC areas for Banded Morwong (Areas 1, 2 and 3) from Low Head on the north coast to Whale Head in the south. Areas 4 and 5 are currently undeveloped.

Current assessment

Biological characteristics

Size composition

Significant changes in the size composition between the late 1990's and the early 2000's raised concerns about the Banded Morwong stock (Fig. 5.2 and Ziegler et al. 2007a). Size composition appears to have stabilised since, with average size of Banded Morwong around 36 cm with fewer large, legal size, fish being present in landings (Fig. 5.2). Comparison between average growth curves show that there are fewer differences between female growth curves post-2007 than between growth curves pre-2007, suggesting that the acceleration of growth (which was evident in the early 2000's) may also have stabilised.

Age composition

Age composition also showed signs of change from the late 1990's (Fig. 5.2 and Ziegler et al. 2007a). The age structure is now dominated by young fish around 4–5 years old (Fig. 5.2). There are relatively very few individuals older than 15 years old in current population samples compared to the late 1990's.




Figure 5.2. Size composition of Banded Morwong by year. n is the sample size, the solid lines represent the minimum and maximum size limits. Year is the year in which the majority of that quota season involved, i.e. 2012 is the 2012 quota season beginning May 2012 and finishing February 2013.




Figure 5.3 Age composition of Banded Morwong by year. n is the sample size. Year is the year in which the majority of that quota season involved, i.e. 2012 is the 2012 quota season beginning May 2012 and finishing February 2013.

Catch, effort and CPUE

The assessment year for Banded Morwong is based on the quota year (1st March to end of February the following year) rather than fishing years (1st July to end June the following year) as for the other scalefish species. The present Banded Morwong assessment includes data up to and including the 2013/14 quota year (which ended February 28, 2014). Note that Banded Morwong catch figures elsewhere in this report are reported for financial years to allow for comparison between species.

Commercial catches under the quota system are reported as numbers of fish rather than weight, which are then converted to weight based on an average of 1.3 kg per fish. State-wide catches have been relatively stable since the introduction of the quota system in 2008/09, and in 2013/14 were estimated at 34.2 t (Fig. 5.2). The total catch in the TAC area (Region 1–3 in Fig.5.1) was 30.8 t (6.4 t in 1 and 14.3 t in 2 and 12.1 t in 3), which represented 94% of the 2013/14 TAC.

While catches in the Region 3 (SEC) have remained relatively stable over time, catches in Region 2 (EC) and Region 1 (NEC) have been steadily declining since 2009/10 (Fig. 5.2 and Fig. 5.4). Reference points pertaining to regional catches have not been triggered in any of the assessment regions (see Reference points section further below).




Figure 5.2 Banded Morwong commercial catches (t). Left: Total state-wide (Total) and graball net (GN) catches, and best estimates of recreational catches (blue squares); Right: regional graball net catches in the TAC areas 1 NEC, 2 EC and 3 SEC. Note NEC catches




Figure 5.3 Left: State-wide commercial effort by method based on gear units and day fished relative to 1995/96. Right: Relative effort in day fished in the TAC areas 1 NEC, 2 EC and 3 SEC.

In 2013/14 effort in both days fished and gear units (100m net hour) decreased compared to 2012/13 (Fig. 5.3). Fishers have progressively reduced their fishing activity and deployed less gear on average for each day fished over the last 10 years (Fig. 5.4). Increasing levels of seal interactions over time has driven affected fishers to fish with less gear or doing fewer sets each day to reduce losses to seals (Ziegler et al. 2006).




Figure 5.4 (A) Banded Morwong catches (tonnes per year) and (B) effort (days per year) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Overall standardised catch rates in the whole TAC area fell between 1995/96 and 1999/2000, before rising back to the 1995/96 levels (Fig. 5.5). Since 2010/11 however, catch rates have sharply declined and are now at approximately just over half of 1995/96 levels. Region 1 (NEC) saw an increase in catch rates between 2009/10 and 2010/11 to levels comparable to the late 1990's, however in 2012/13 catch rates declined again before rising sharply in 2013/14. Region 3 (SEC) also saw an increase in catch rates between 2009/10 and 2011/12 before catch rates fell in 2012/13 to their 2nd lowest level since 1995/96. In 2013/14 catch rates have rebounded in Region 3 (SEC) similar to Region 1 (NEC). Catch rates in Region 2 (EC) remained relatively stable from 2000–2010 before declining to record lows. As a consequence, catch rates were below the reference points for Region 2 (EC), Region 3 (SEC) and for the whole State (see reference point section further below).




Figure 5.5 Banded Morwong standardised grabball catch per unit effort (CPUE by days fished) relative to 1995/96, from the whole TAC area (top left), and in the TAC areas 1 NEC, 2 EC and 3 SEC (bottom left and right).

Selected stock assessment results

Details of the Banded Morwong model can be found in Ziegler et al. (2007b). The model was used to project mature biomass and CPUE into the future under various TAC scenarios. These included the 2012/13 TAC (38.8 t), the 2013/14 TAC (37.2 t), the 2014/15 TAC (35.7 t) and a comparative TAC of 30 t. State-wide and for all regions, only the modelled TAC of 30 t achieves a reversal of the decreasing trend in mature biomass over the next 10 years, bringing the 2021 mature biomass back to the 2013 level (Fig. 5.6).




Figure 5.6 Predicted mature biomass from the Banded Morwong model under the 2012/13 (38.8 t), 2013/14 (37.2 t), 2014/15 (35.7 t) TAC settings and a comparative 30 t TAC for the whole State, Region 2 (East) and Region 3 (South-east).

Limit reference points under the various TAC were calculated using the model, namely a 90% probability of the mature biomass catch limit being above 30% of the virgin biomass by 2016 and a 90% probability of the catch per unit effort being above the 10-year low (2000/01) by 2016. Region 3 (SEC) and the State as a whole had greater than a 90% probability of reaching the mature biomass reference point by 2016 under all TAC's, and Region 3 (SEC) had greater than a 90% probability of reaching the CPUE reference point by 2016 under the 2014/15 TAC and the 30t TAC. Region 2 was not predicted to reach the limit reference points regardless of the TAC applied (Table 5.1).

Table 5.1 Probabilities of the mature biomass being above 30% of the virgin biomass (with a 90% probability) by 2016 and the CPUE being above the 10-year low by 2016 under various TAC scenarios.

Region	Limit reference point	TAC 2012/13 (38.8t)	TAC 2013/14 (37.2t)	TAC 2014/15 (35.7t)	TAC 30t
2- EC	CPUE	89	95	95	100
	Mature biomass	18	18	27	63
3- SEC	CPUE	68	85	92	99
	Mature biomass	100	100	100	100
Statewide	Mature biomass	94	96	96	99

Reference points

The 2014/15 TAC predictions were used to assess the limit reference points for mature biomass and catch per unit effort.

Performance indicators	Reference points	Breached?	By how much?
Fishing mortality	• Catch region 1 >30% TAC (>10.7 t)	No	
	• Catch region 2 > 65% TAC (>23.2 t)	No	
	• Catch region 3 > 40% TAC (>14.3 t)	No	
	• Outside TAC area > 10 t (>3.6 t)	No	
	• Commercial catch <90% TAC (<32.1 t)	No	
Biomass	• Mature biomass catch limit >30% of the virgin biomass with a 90% probability in 2016	2- EC	Yes
		3- SEC	No
		State	Yes
	• Catch per Unit Effort > the 10-year low (2000/01) with a 90% probability in 2016	2- EC	Yes
		3- SEC	No
	• Catch rate <0.9 x average from reference period 2000/01 to 2006/07	1- NEC	No
		2- EC	Yes
3- SEC		Yes	28%
State		Yes	32%
Stock stress	• Significant change in the size/age composition	Yes	Change size/age composition, acceleration of growth, earlier maturity (Ziegler et al. 2008), appears to have recently stabilised

	<ul style="list-style-type: none"> Change in catches of non-commercial fish relative to 1990/91 to 1997/98 or high incidental/undersize mortality 	No	
	<ul style="list-style-type: none"> Significant numbers of unhealthy fish landed 	No	

Stock status

TRANSITIONAL DEPLETING

While none of the fishing mortality indicators were exceeded, several of the biomass reference points were breached. Region 2 (EC) appears particularly vulnerable, breaching both the mature biomass and catch-per-unit effort reference points. Even under a 30 t TAC (corresponding to 5.7 t less than the current 2014/15 TAC), the mature biomass for Region 2 is not predicted to reach acceptable limits by 2016.

Overall, the model estimated a decrease in mature biomass between 1990-2009 in Region 2 (EC) from around 436 t to 134 t or 31% of the initial levels, and in region 3 (SEC) from around 218 t to 97 t or 44% of the initial levels. These estimates change from one assessment to the next. In this case, the substantial (and unexpected) decline in catch-rate over the last two years has changed model results substantially, providing a reduction in both the estimate of the initial (1990) stock biomass and the current stock biomass.

The structure of the fished population has changed substantially in both regions. Recent stock rebuilding in the southern stocks was based on strong recruitment pulses in the 2000s coupled with higher productivity through increased growth rates and earlier maturation (Ziegler et al. 2007a). As a result, the fishery in both regions is now largely recruitment-driven with only a small proportion of the catch made up by older females. With continually high harvest rates above the internationally recognised reference points for mature biomass of $H_{40\%}$ or $H_{30\%}$ in recent years, the fishery mainly removes and depends on newly-recruited fish from the populations at a potentially unsustainable rate. While a number of year classes contribute to the fishery, even a relatively short period of low recruitment could lead to substantial declines in catch rates and mature biomass.

Banded Morwong have a particularly long larval phase, which may decrease the likelihood of successful recruitment as climate driven changes to oceanography (sub-Antarctic convergence and east Australian current) disperse larvae further from spawning grounds. This occurrence has recently been identified as a major factor causing poor recruitment in east Australian Jackass Morwong stocks and has led to a significant decline in the productivity of the fishery (Wayte 2013).

The higher dependence on new recruitment in both regions was reflected in the risk assessment. With the current TAC the model run in 2013 predicted that mature biomass and catch rate would continue to decline. With the scheduled TAC for 2014/15 the model predicts that catch rate and mature biomass will decrease before stabilising. Model based reference points for CPUE on the East Coast are met under the 2014/15 TAC cut, however even a reduction to 30 t would be insufficient to meet the reference points for mature biomass on the East Coast or Statewide. Given uncertainty in the assessment and the magnitude of inter-annual fluctuations in CPUE the time-horizon for the current reference points (2016) may be too short and may suggest un-necessarily overly conservative management action in response to these fluctuations.

Due to the biological characteristics of Banded Morwong, stock recovery under all considered TACs is expected to be a slow process. There is considerable uncertainty in the assessment about the basic population dynamics of Banded Morwong, mainly in regards to the potential separation of stocks in fished onshore and unfished offshore populations. In addition, the

species' productivity appears to be able to respond strongly and quickly to stock biomass changes.

The Banded Morwong TAC is set as a tonnage but implemented in numbers. Each fisher is allocated a certain number of fish by dividing their allocated catch (in kg) by a predetermined average fish weight. The same average weight is used throughout the state. This is an unusual strategy resulting from practical difficulties in weighing live fish. There are indications from both catch records and anecdotal reports that this system can lead to high grading of small fish. The motivation is that by discarding smaller fish, fishers can increase their average fish weight and thereby increase their revenue. Potential problems include:

- Exceeding the TAC – An increase in average weight above the assumed value will result in a catch exceeding the TAC.
- Preferential targeting of areas with large fish – Average catch size can also be boosted by targeting areas with large fish. The current system provides an economic incentive to do so.
- Biological effects of a changed size structure – High grading will change the size structure, the biological effects of this have not been quantified.
- Difficulties in interpreting CPUE data – High grading (by discarding small fish) lowers CPUE.

DPIPWE is proposing to move the Banded Morwong Fishery to a weight based system in the remake of the Scalefish Management Plan scheduled for public consultation in 2015.

6. Barracouta

Thyrsites atun

STOCK STATUS	Not assessed
Large fishery for Barracouta in the 1960's but little market for the species nowadays.	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends



Thyrsites atun
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source								
Habitat	<ul style="list-style-type: none"> Open water and coastal bays. Down to 550 m depth. 	Edgar (2008) Kailola et al. (1993)								
Distribution	<ul style="list-style-type: none"> From mid-Western Australia to south Queensland, and around Tasmania. Also widely in the southern hemisphere in temperate latitudes. 	Edgar (2008)								
Diet	<ul style="list-style-type: none"> Pelagic crustaceans, cephalopods, fishes (e.g. anchovy and Jack Mackerel) 	Nakamura and Parin (1993)								
Movement and stock structure	<ul style="list-style-type: none"> Schooling fish Some stocks undertake annual migrations lasting 6–9 months and covering several hundreds of km. Also moves through the water column from 200 m depth to the surface At least 5 stocks: 3 in south-eastern waters, 1 in South Australia and 1 in Western Australia 	Paul (2000) Kailola et al. (1993) Blackburn and Gartner (1954)								
Natural mortality	<ul style="list-style-type: none"> $M = 0.3$ 	Hurst et al. (2012)								
Maximum age	<ul style="list-style-type: none"> At least 10 years, potentially up to 15 years. 	Kailola et al. (1993) Hurst et al. (2012)								
Growth	<ul style="list-style-type: none"> Maximum length: 1.4 m Maximum weight: 6 kg Growth described by von Bertalanffy growth function $L = L_{\infty}(1 - e^{-k(t-t_0)})$ where L is the fork length (cm), t is the age (years), L_{∞} is the average maximum length for the species, k is a constant and t_0 is the (theoretical) age where length equals zero. Parameters estimates are : <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Sex</th> <th>L_{∞}</th> <th>k</th> <th>$t_{0\infty}$</th> </tr> </thead> <tbody> <tr> <td>Combined</td> <td>91</td> <td>0.42</td> <td>-0.25</td> </tr> </tbody> </table> Length-weight relationship was estimated at $W = 0.1064 L^{2.2385}$ for females and males combined where W is weight (g) and L is the fork length (cm). 	Sex	L_{∞}	k	$t_{0\infty}$	Combined	91	0.42	-0.25	Edgar (2008) Nakamura and Parin (1993) Grant et al. (1978) Blackburn (1960)
Sex	L_{∞}	k	$t_{0\infty}$							
Combined	91	0.42	-0.25							
Maturity	<ul style="list-style-type: none"> Sexual maturity at about 50–60 cm FL and about 2–3 	Hurst et al. (2012)								

	years of age	
Spawning	<ul style="list-style-type: none"> October to March in Tasmania 	Kailola et al. (1993)
Early life history	<ul style="list-style-type: none"> Little data. Eggs are pelagic and juveniles inhabit sheltered waters of southern bays and estuaries. 	Kailola et al. (1993) Hurst et al. (2012)
Gillnet post release survival	<ul style="list-style-type: none"> NA 	

Background

Barracouta used to be the subject of a large commercial trolling fishery in Tasmania in the 1960's and 1970's, with catches ranging from 600 – 1 600 t (Kailola et al. 1993). Demand for Barracouta, however, declined in the mid-1970's and there is now little commercial fishing for the species. Barracouta abundance also fluctuates greatly in State waters annually.

FISHING METHODS	Mostly troll and handline
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence) Output control: <ul style="list-style-type: none"> Possession limit of 15 individuals for recreational
MAIN MARKET	Local

Current assessment

Catch, effort and CPUE

Trolling is the main fishing method used to target Barracouta. Catches peaked in the early 2000's with maximum of 132.1 t but gradually declined from 2004/05 to 1.1 t in 2013/14 (Fig. 6.1A). Catches and fishing effort were concentrated in southern Tasmania during the past five years; however the 2013/14 catch and effort were predominantly off Devonport (Fig. 6.2).

After the peak in the early 2000's, effort has declined and stabilised since 2007/08 at a low level (Fig. 6.1B). Catch rates have been relatively stable over time but have been at their lowest levels in 2013/14 (Fig. 6.1C); however, it is likely that fishers utilising fishing gears historically used to target Barracouta are now targeting other species and as a result, catch rate is unlikely to be a reliable indicator of abundance.

Barracouta are targeted and also taken as by-product by the recreational sector. Catches were estimated at 46.9 t in 2000/01 (Lyle 2005), 10.8 t in 2007/08 (Lyle et al. 2009) and 31 t in 2012/13 (Lyle et al. 2014b), the latest estimate being considerably greater than the commercial harvest in recent years.

Barracouta





Figure 6.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. TR= troll, HL= hand line, SEC= south-east coast, EC= east-coast.

A) Catch



B) Effort




Figure 6.2 (A) Barracouta catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (351.5 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (19.3 t)	Yes	18.2 t (94.3%)
	• Catch increases by > 30% from previous year (>1.48 t)	No	
	• Catch decreases by > 30% from previous year (<0.80 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>773 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0145 t/days fished)	Yes	0.008 t/days fished (54.5%)

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (85.2 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (25.0 t)	Yes	23.9 t (95.6%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period (46.9 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (43.8% in 2007/08)	Yes	Latest estimate (2012/13): 96.6%
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0085)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status


NOT ASSESSED

Historically, the population of Barracouta has undergone large fluctuations in population size and availability, possibly linked to recruit variability and environmental factors. Catches of Barracouta have been declining steadily since the mid-2000s due to a decrease in targeted effort as a result of a lack of market opportunities. The increase in recreational catch proportion mainly reflects the sharp fall in commercial landings rather than increased recreational catch. Consequently, catches and catch rates are not considered indicative of stock status. As such, the status of the stock is uncertain and substantial effort would be required to gather sufficient data to assess it.

7. Bastard Trumpeter

Latridopsis forsteri

STOCK STATUS	TRANSITIONAL DEPLETING
This species is mainly taken as a by-product rather than a target species, and the recreational catch exceeds commercial landings.	
IMPORTANCE	Key
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends



Latridopsis forsteri
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source								
Habitat	<ul style="list-style-type: none"> Exposed reefs and sandy bottom down to 160 m depth. 	May & Maxwell (1986) Edgar (2008)								
Distribution	<ul style="list-style-type: none"> Sydney (New South Wales) to southern South Australia, Tasmania, southern New Zealand 	Edgar (2008)								
Diet	<ul style="list-style-type: none"> Small fish, invertebrates. 	Edgar (1997)								
Movement and stock structure	<ul style="list-style-type: none"> Schooling fish, usually in small numbers. Large individuals occur in deeper waters. Juveniles tend to remain associated with areas of reef for periods of time No information on the stock structure. 	Edgar et al. (2004) Gomon et al.(2008)								
Natural mortality	<ul style="list-style-type: none"> Undetermined. 									
Maximum age	<ul style="list-style-type: none"> Up to 20 years 	Murphy and Lyle (1999)								
Growth	<ul style="list-style-type: none"> Maximum length: 65 cm Maximum weight: 4 kg Growth described by von Bertalanffy growth function $L = L_{\infty}(1 - e^{-k(t-t_0)})$ where L is the fork length (cm), t is the age (years), L_{∞} is the average maximum length for the species, k is a constant and t_0 is the (theoretical) age where length equals zero. Parameters estimates are : <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Sex</th> <th>L_{∞}</th> <th>k</th> <th>$t_{0\infty}$</th> </tr> </thead> <tbody> <tr> <td>Combined</td> <td>59.4</td> <td>0.144</td> <td>-2.9</td> </tr> </tbody> </table> 	Sex	L_{∞}	k	$t_{0\infty}$	Combined	59.4	0.144	-2.9	Edgar (2008) Gomon et al.(2008) Murphy and Lyle (1999)
Sex	L_{∞}	k	$t_{0\infty}$							
Combined	59.4	0.144	-2.9							
Maturity	<ul style="list-style-type: none"> Sexual maturity at sizes >45 cm and ages > 4 years. 	Murphy and Lyle (1999)								

Spawning	<ul style="list-style-type: none"> Believed to spawn in late winter 	Murphy and Lyle (1999)
Early life history	<ul style="list-style-type: none"> Small juveniles settle from the plankton on reefs in large numbers at intervals of several years. 	Edgar (2008)
Recruitment	<ul style="list-style-type: none"> Variable. No-stock recruitment relationship established. 	
Gillnet post release survival	<ul style="list-style-type: none"> High: 95 – 83% depending on gillnet soak duration 	Lyle et al (2014a)

Background

Bastard Trumpeter was one of the first fish species to have been commercially exploited in Tasmania. Their apparent abundance around reefs close to newly established Hobart meant that they were an important source of seafood for the fledgling colony. Their exploitation was further aided by the relative ease at which they could be caught using gillnets set within accessible shallow inshore reefs.

Bastard Trumpeter are taken today almost exclusively by graball nets. Bastard Trumpeter have also long been recognised as an important fish for recreational fishers. The species resides on inshore reefs until about 4–5 years of age (and approximately 50 cm long) before moving offshore into deeper water as they approach maturity, apparently remaining in that habitat for the remainder of their lives (Harries and Lake 1985, Murphy and Lyle 1999). Hence, both commercial and recreational fisheries are based almost entirely on juvenile fish.

FISHING METHODS	Graball net
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence) Output control: <ul style="list-style-type: none"> Possession limit of 10 individuals for recreational Trip limit 200 kg for commercial Legal minimum length (380 mm TL)
MAIN MARKET	Local

Current assessment

Catch, effort and CPUE

Bastard Trumpeter catches have been declining steadily since the mid-1990's and reached a historic low of 8.2 t in 2013/14 (Fig. 7.1A). Bastard Trumpeter are taken almost exclusively by graball net from inshore waters off the east, south and west coasts (Fig. 7.2). Bastard Trumpeter have predominantly been taken by recreational fishers in recent years, although the estimated catch in 2012/13 was also a historic low of 9.8 t (Lyle et al. 2014b) and is considerably lower than the 2010 estimate of 27.3 t (Lyle and Tracey 2012).

Graball effort has followed a similar downward trend to catches since the mid-1990's (Fig. 7.1B). Daily catch rates have remained relatively stable over time with a slight decrease since 2006/07 (Fig. 7.1C). This lack of obvious trend, despite the sharp decrease in catches, presumably reflects the fact that Bastard Trumpeter are taken primarily taken as by-product rather than as a target species. This is exacerbated by the fact that the majority of gillnetting effort now targets Banded Morwong with 140 mm gillnets, which only select the largest Bastard Trumpeter. Previously, a larger proportion of fishers used smaller mesh sizes (~114 mm) to target Bastard Trumpeter and/or Blue Warehou.

Bastard trumpeter





Figure 7.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and days fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. GN= graball net, EC= east coast, SEC= south-east coast and WC= west coast.

A) Catch



B) Effort




Figure 7.2 (A) Bastard Trumpeter catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (63.3 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (34 t)	Yes	25.8 t (75.9%)
	• Catch increases by > 30% from previous year (>12.2 t)	No	
	• Catch decreases by > 30% from previous year (<6.6 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>2717 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0091 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (47.5 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (21.0 t)	Yes	12.7 t (60.6%)
	• Catch variation from the previous year above the greatest inter-annual increase from the reference period (7 t)	No	
	• Catch variation from the previous year above the greatest inter-annual decrease from the reference period (-11.4 t)	No	
	• Latest recreational catch estimate > recreational catch estimate from the reference period (24 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (73.6% in 2010)	No	
Biomass	• CPUE < 3 rd lowest CPUE value from the reference period (0.0106 t/days fished)	Yes	0.0005 t/day fished (4.7%)
	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0011)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

TRANSITIONAL DEPLETING

Due to the very low landings in 2013/14, the lowest catch and fishing mortality reference points were breached. As Bastard Trumpeter is a by-product species, total catch rather than catch rate may be a better indicator of abundance/availability for the species. As such, the trend in commercial production suggests that current inshore populations are at historically low levels.

In accordance with this observation, industry and recreational representatives have expressed concerns about the scarcity of the species in recent years, although a lack of market demand for Bastard Trumpeter appears to be a factor influencing catches. Onboard observation suggests that legal sized Bastard Trumpeter are sometimes discarded by Banded Morwong fishers and given that the majority of gillnetting effort is now targeted at Banded Morwong using larger gillnets than those used historically to target Bastard Trumpeter, both catch and catch rate may be inaccurate. Nevertheless, fishing practices are likely to have remained fairly consistent in recent years (2007/08 – present) so the low, stable catch rate trend is likely to be representative of a population that has not substantially rebuilt despite significant reduction in both commercial and recreational gillnet effort.

The Tasmanian fishery is based almost entirely on juveniles. As fish grow, they appear to move offshore and are rarely caught. No information is available on the adult portion of the population but it is clear that fishing pressure exerted on those individuals that evade the inshore fishery is very low (by-catch in shark nets, trawl, Danish seine or deep-water fish traps used by the Commonwealth SESSF appears to be negligible). The species exhibits high recruitment variability resulting in short-term variation in catches, which has been a feature of this fishery over the past century (Harries and Croome 1989). Anecdotal reports and low inshore catches suggest that recruitment level has been low in recent years. Since commercial and recreational fisheries are based on juveniles, recruitment as well as growth overfishing is likely to be occurring.


It is worth noting that the stabilisation of the catch since 2009/10 corresponds to the introduction of several management measures for the species (increase in the minimum legal size, introduction of commercial and recreational catch limits). However, the current minimum size limit of 38 cm TL is still well below the size at maturity (>45 cm FL, Murphy and Lyle 1999). Increasing the minimum size limit to above the size at maturity would be beneficial to the stock but would also effectively close down the current commercial and recreational fisheries for the species.

Given the continued low catch rate, the lack of recruitment and the fact that the fishery predominantly targets juveniles, Bastard Trumpeter are classified as transitional depleting.

8. Blue Warehouse

Seriolella brama

STOCK STATUS	OVERFISHED
Commonwealth-managed species (See: http://www.afma.gov.au/portfolio-item/blue-warehouse/).	
Sporadically abundant in Tasmanian waters. Despite a reduction in Total Allowable Catch (TAC) for the Commonwealth fishery, stocks have shown little signs of recovery.	
IMPORTANCE	Key
STOCK(S)	Tasmanian Scalefish Fishery/Southern and Eastern Scalefish and Shark Fishery (Commonwealth)
INDICATOR(S)	Catch, effort and CPUE trends



Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> Inshore reefs/harbours and open water down to 400 m depth. 	Edgar (2008) Smith (1994)
Distribution	<ul style="list-style-type: none"> New South Wales to South Australia, Tasmania, New Zealand 	Edgar (2008) Gomon et al.(2008)
Diet	<ul style="list-style-type: none"> Invertebrates (mainly salps), krill, crabs, squids 	Gavrilov and Markina (1979) Annala (1994) Bulman et al. (2001)
Movement and stock structure	<ul style="list-style-type: none"> Schooling fish, highly mobile. Small juveniles pelagic, commonly in association with jellyfish in open coastal waters. Sub-adults often found in sheltered waters of large marine embayments. Although genetics has not confirmed separate stocks, there are indications of population structuring. It is likely that two stocks occur in southern Australian waters (east and west of Bass Strait). 	Gavrilov and Markina (1979) AFMA (2008, 2011) Bruce et al. (2001b) Robinson et al. (2008)
Natural mortality	<ul style="list-style-type: none"> Estimated between $M=0.30$ and $M=0.45$. 	Knuckey and Sivakumaran (2001)
Maximum age	<ul style="list-style-type: none"> Up to 15 years 	AFMA (2012)
Growth	<ul style="list-style-type: none"> Maximum length: 90 cm Maximum weight: 4 kg Growth (in New Zealand) described by von Bertalanffy growth function $L = L_{\infty}(1 - e^{-k(t-t_0)})$ 	Gomon et al.(2008) Frimodt (1995) BWAG (1998)

	<p>where L is the length (cm), t is the age (years), L_{∞} is the average maximum length for the species, k is a constant and t_0 is the (theoretical) age where length equals zero.</p> <p>Parameters estimates are:</p> <table border="1"> <thead> <tr> <th>Sex</th> <th>L_{∞}</th> <th>k</th> <th>$t_{0\infty}$</th> </tr> </thead> <tbody> <tr> <td>Combined</td> <td>54.65</td> <td>0.37</td> <td>-0.67</td> </tr> </tbody> </table> <ul style="list-style-type: none"> Length-weight relationship was estimated at $W = 0.03L^{2.9}$ for females and males combined, where W is weight (g) and L is the total length (cm). 	Sex	L_{∞}	k	$t_{0\infty}$	Combined	54.65	0.37	-0.67	
Sex	L_{∞}	k	$t_{0\infty}$							
Combined	54.65	0.37	-0.67							
Maturity	<ul style="list-style-type: none"> Size-at-50% maturity estimated at 36 cm (3.67 years) for females. Batch fecundity (BF): $\ln(BF) = 2.614\ln(L) + 2.366$, where L is length in cm. 	Knuckey and Sivakumaran (2001)								
Spawning	<ul style="list-style-type: none"> Peak spawning in winter, with major regional differences in the magnitude and timing of spawning. Major spawning ground on the central-west and north-west coasts in Tasmania 	Bruce et al. (2001b)								
Early life history	<ul style="list-style-type: none"> Larvae restricted to shelf and slope waters. Larvae likely to be transported by Zeehan Current from spawning grounds of western Tasmania to south-eastern Tasmania nursery areas. Larvae settle to the bottom at length > 14.5 mm BL 	Bruce et al. (2001b) Neira et al. (1998)								
Recruitment	<ul style="list-style-type: none"> Variable. No-stock recruitment relationship established. 									
Gillnet post release survival	<ul style="list-style-type: none"> Low: 35%, but small legal minimum length so rarely discarded. 									

Background

Blue-warehou occur seasonally in Tasmanian inshore waters, the region representing the southern-most extent of the species' distribution. The availability of Blue Warehou in coastal waters is assumed to be influenced by prevailing oceanographic conditions and availability of prey species, in particular salps. These factors produce marked inter-annual variability in abundance and hence catch taken from State waters.

Blue Warehou is a Commonwealth managed species and a Memorandum of Understanding (MoU) exists to cover catches from Tasmanian State Fishing Waters. Within the context of this MoU, State catches of Blue Warehou are to be managed within historical levels. The species is assessed at the Commonwealth level with Tasmanian (and other state) catches included in the stock assessment modelling.

FISHING METHODS	Mainly graball net, also small mesh nets and seine net
MANAGEMENT METHODS	<p>Input control:</p> <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence) <p>Output control:</p> <ul style="list-style-type: none"> Possession limit of 20 individuals for recreational fishers Minimum size (250 mm TL)
MAIN MARKET	Local and interstate

Current assessment

Catch, effort and CPUE

In Tasmania, Blue Warehou is taken primarily in graball nets with lesser quantities taken by small mesh nets and seine (Fig. 8.1A). They are captured using a variety of methods by Commonwealth fisheries including other gillnet categories (e.g. shark gillnets), Danish seine and trawl.

Due to low availability since the early 2000's, the species has been rarely targeted. The current catch of 5.2 t is down from the previous year (8.5 t) and represents the historic minimum since records have been maintained. Peak Tasmanian landings were 317.6 t in 1991/92, which corresponded with the peak Australia wide landings of almost 3,000 t (AFMA 2012). Two stocks of Blue Warehou occur in southern Australian waters, the east and the west Bass Strait stocks (Bruce et al. 2001b). The Tasmanian fishery is mainly centred off the south-east coast and thus probably targets the eastern stock (Fig. 8.2). Catches are also taken off the north-west coast, probably involving the western stock.

Blue Warehou are also targeted by recreational fishers using gillnets, and to a lesser extent line fishing. Recreational catches were historically lower than commercial catches (Fig. 8.1A), although the estimates in 2010 were similar catches for both sectors (32.5 t for recreational and 37.5 t for commercial) and the 2012/13 recreational catch of 15.4 t (Lyle et al. 2014b) exceeded the commercial catch of 8.5 t.

Blue warehou





Figure 8.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. GN= graball net, SEC= south-east coast, EC= east coast.

Following an increase in commercial graball effort between 1995/96 and 1998/99 that resulted in increased catches, effort has fallen to a substantially lower level (Fig. 8.1B). Current effort is amongst the lowest since 1995/96, which is largely in response to the reduced availability of Blue Warehou in Tasmanian waters but has also been influenced by the quota management system introduced for the Banded Morwong sector.

Graball catch rates increased markedly in the latter part of the 1990s reflecting increased availability, and targeting, of Blue Warehou around Tasmania at the time (Fig. 8.1C). Catch rates have since declined, but remained relatively stable throughout the 2000's before a further decline during the last three years.

A) Catch



B) Effort




Figure 8.2 (A) Blue Warehouse catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (317.6 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (82.3 t)	Yes	77.1 t (93.7%)
	• Catch increases by > 30% from previous year (>11 t)	No	
	• Catch decreases by > 30% from previous year (<5.9 t)	Yes	0.7 t (11.8%)
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (2215 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (0.0183 t/days fished)	Yes	0.0032 t/days fished (17.5%)

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Commercial catch limit of 318 t as per Memorandum of Understanding (MoU)	No	
	• Catch > 3 rd highest catch value from the reference period (186.1 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (27.5 t)	Yes	22.3 t (81%)
	• Catch variation from the previous year above the greatest inter-annual increase from the reference period (84.7 t)	No	
	• Catch variation from the previous year above the greatest inter-annual decrease from the reference period (151.6 t)	No	
	• Latest recreational catch estimate > recreational catch estimate from the reference period (65.3 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (57.3%, in 2010)	Yes	Latest estimate (2012/13): 63.6%
Biomass	• CPUE < 3 rd lowest CPUE value from the reference period (0.0229 t/days fished)	Yes	0.0077 t/days fished (33.7%)
	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0126)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

OVERFISHED

The availability of Blue Warehou in Tasmanian inshore waters is influenced by a range of environmental factors as well as stock size. As such, despite several reference limits being

breached, catch and CPUE may not be a good indicator of abundance/availability for the species.

Nevertheless, decreasing catch and catch rates over the last 20 years are almost certainly linked to reduced biomass. This is predominantly a result of overfishing by Commonwealth and State fisheries during the 1990's when >2500 t were caught in several years and >1000 t annually between 1987 and 1998 (AFMA 2012). These figures include all state landings and the Tasmanian state landings accounted for around 10% of the total Australian Blue Warehou landings throughout much of this period (AFMA (2012); Appendix 3).

The Commonwealth Total Allowable Catch (TAC) for Blue Warehou has been progressively decreased since 2003, and was further reduced to 118 t for 2012/13, 2013/14 and 2014/15 since both the western and eastern stocks were considered to be overfished and unlikely to recover under the existing management arrangements (AFMA 2012). Recent Commonwealth catches have been below the TAC suggesting there is little, or no, targeting occurring and there is little benefit in further reductions. Further, Commonwealth vessels are not permitted to fish in two known Blue Warehou spawning grounds off western Victoria and Tasmania during their spawning season.


Despite the Commonwealth management measures outlined above, there have been few signs of recovery for the species and the Fishery Status Reports 2013-14 classified the status of Blue Warehou stocks as overfished (for biomass) and uncertain (for fishing mortality) (Georgeson et al. 2014). Blue Warehou were not assessed in the 2014 Status of Australian Fish Stocks but the most recent Blue Warehou Rebuilding Strategy suggests there has yet to be a significant recovery (AFMA 2014). As such, Blue Warehou is assessed as overfished in Tasmanian waters.

9. Flathead

Tiger Flathead *Platycephalus richardsoni*

Southern Sand Flathead *Platycephalus bassensis*

STOCK STATUS	SUSTAINABLE (Tiger Flathead) NOT ASSESSED (Southern Sand Flathead)
Tiger flathead is also a Commonwealth-managed species (See: http://www.afma.gov.au/portfolio-item/tiger-flathead/) Both Flathead species are caught by commercial and recreational fishers in Tasmania, however, Tiger Flathead are the dominant species in the commercial catch whereas Sand Flathead dominates the recreational catch.	
IMPORTANCE	Key
STOCK(S)	Tasmanian Scalefish Fishery/ Southern and Eastern Scalefish and Shark Fishery (Commonwealth)
INDICATOR(S)	Catch, effort and CPUE trends



Species biology

Parameters	Estimates	Source
Habitat	<u>Tiger</u> : Exposed sand and silt habitat. 10-400 m depth <u>Sand</u> : Sheltered sand and silt habitat. Down to 100m depth.	Edgar (2008) Tilzey et al. (1990)
Distribution	<u>Tiger</u> : Victoria to New South Wales and around Tasmania. <u>Sand</u> : From Western Australia to New South Wales and around Tasmania.	Edgar (2008) Gomon et al.(2008)
Diet	<u>Tiger</u> : Fish <u>Sand</u> : Fish and crustaceans	Coleman and Mobley (1984) Ayling et al. (1975)
Movement and stock structure	<u>Tiger</u> : Young inhabit shallow waters of the continental shelf and move into the outer shelf zone as they reach maturity <u>Sand</u> : Seasonal movements between inshore and offshore in east and south-east Tasmania	Kailola et al. (1993) Jordan 1998
Natural mortality	<u>Tiger</u> : <i>M</i> between 0.21 and 0.46 <u>Sand</u> : <i>M</i> between 0.28 and 0.59 (per year)	Klaer (2010) Bani (2005)
Maximum age	<u>Tiger</u> : 12 years <u>Sand</u> : 20 years	Rowling (1994) Bani (2005)

	species	
Gillnet post release survival	<ul style="list-style-type: none"> Moderate: 50% (all soak durations and including both graball and mullet nets) 	Lyle et al. (2014a)
Rod and line post release survival	<ul style="list-style-type: none"> High: >99% for circle hooks and 94–97% for conventional hooks. 	(2007)

Background

Several species of Flathead occur in Tasmanian waters, but the commercial catches are dominated by Tiger Flathead taken by Danish seine. Sand Flathead are caught to a lesser extent by handline and as a by-product in graball nets. However, the two species were not routinely distinguished in catch returns until 2007; consequently, catches prior to 2007 have been inferred based on fishing gear.

Sand Flathead are an important recreational species, which is targeted by handline or rod and reel. Tiger Flathead only constitute a minor component of the recreational harvest (around 10%).

FISHING METHODS	Danish seine (Tiger Flathead), hand line, rod and reel, and graball net (Sand Flathead)
MANAGEMENT METHODS	<p>Input control:</p> <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence) <p>Output control:</p> <ul style="list-style-type: none"> Possession limit of 30 individuals (all Flathead species combined) for recreational Minimum size (300 mm TL)
MAIN MARKET	Mostly local

Current assessment

Catch, effort and CPUE

The 2013/14 total commercial catch for flathead species is 31.1 t (20.2 t Tiger Flathead, 6.8 t Sand Flathead and 4.1 t unspecified Flathead). The commercial fishery for Flathead has not undergone major changes in its operations since 1995/96. It was therefore possible to back calculate catches for each species prior to 2007, when species was not distinguished, based on the average proportion of species by gear type from 2007/08 to 2011/12 (Fig. 9.1). Tiger Flathead catches declined steadily between 2000/01 and 2003/04 but more than doubled in 2005/06 (Fig. 9.1A and 9.2A). Since then catches have fluctuated between 30 t and 70 t per annum, but are currently at their lowest level in eight years at 20.2 t. Sand Flathead catches remained relatively stable between 10 and 15 t per annum until 2008/09 but have subsequently declined, reaching 6.8 t in 2013/14 (Fig. 9.1B).




Figure 9.1 Back-calculated annual commercial catch (t) by gear for A) Tiger Flathead and B) Sand Flathead.

Catches in 2013/14 were derived mainly from the south-east and east coasts, with smaller quantities also taken from the north-east (Flinders Island) and north-west coasts (Fig. 9.2A and 9.3). Catch and effort for the latter two regions have declined compared to the previous 5 year period (Fig. 9.3). However, Danish seine catch and effort can be influenced by the species being targeted and an increase in the landings of School Whiting in 2013/14 (See Chapter 18) suggests that lesser effort was directed at Tiger Flathead.

Recreational Flathead catches were estimated at 361 t in 2000/01 (Lyle 2005), 292 t in 2007/08 (Lyle et al. 2009) and 235.9 t in 2012/13 (Lyle et al. 2014b). Sand Flathead are by far the dominant species taken (>90% in numbers) (Fig. 9.2A).

Flathead





Figure 9.2 A) Combined annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Combined commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Combined commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. DS= Danish seine, HL= hand line, SEC= south-east coast, EC= east coast.

Danish seine fishing effort has fluctuated without obvious trend since the mid-1990s, whereas handline effort has declined since 2008/09 (Fig. 9.2B). There have been reports from the commercial sector that Sand Flathead availability has been declining since 2009, which could account for this trend, although hand line catch rates have remained stable over time. Danish seine catch rates have fluctuated strongly in recent years and presumably reflect the degree of targeting of the species (Fig. 9.2C). Peaks in Danish seine CPUE are influenced by a small number of operators that exclusively targeted Tiger Flathead during those years.

A) Catch



B) Effort




Figure 9.3 (A) Combined Flathead catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (165.3 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (51.8 t)	Yes	20.7 t (40%)
	• Catch increases by > 30% from previous year (> 52.2 t)	No	
	• Catch decreases by > 30% from previous year (<28.1 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>1341 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0104 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (63.1 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (50.5 t)	Yes	19.4 t (38.4%)
	• Catch variation from the previous year above the greatest inter-annual increase from the reference period (-32.6 t)	No	
	• Catch variation from the previous year above the greatest inter-annual decrease from the reference period (43.5 t)	No	
	• Latest recreational catch estimate > recreational catch estimate from the reference period (361 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (80.0% in 2007/08)	Yes	Latest estimate (2012/13) 85.5%
Biomass	• CPUE < 3 rd lowest CPUE value from the reference period (0.0130 t/days fished)	Yes	0.0003 t/days fished (2.3%)
	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0020)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status	Tiger Flathead	SUSTAINABLE
	Sand Flathead	NOT ASSESSED

Danish seine catches are highly variable and tend to be inversely related with catches of School Whiting (refer Fig. 2.1), which is also targeted using Danish seine. The 3rd lowest catch reference point was breached, linked to decreasing Sand Flathead catches and low Tiger

Flathead catches. The 3rd lowest CPUE reference point was also breached, although only by a small margin. While commercial catches of Tiger Flathead have been maintained at higher levels in the past, there are additional and significant trawl catches of Flathead (almost exclusively Tiger Flathead) taken from Commonwealth waters as part of the South East Trawl Fishery. Tasmanian catches only constitute a fraction of the overall catches from this stock (extending from New South Wales to Victoria), which has been classified as sustainably fished in recent years (Flood et al. 2012, Woodhams et al. 2013, Flood et al. 2014). In accordance with the Commonwealth assessment, Tiger Flathead is assessed as sustainable in Tasmanian waters. Despite an overall reduction in recreational catches, the decline in commercial catches still resulted in the recreational catch proportion performance indicator being breached, further highlighting the significance of the recreational sector.


Due to a lack of commercial targeting of Sand Flathead their stock status is not known, though undersized fish appear to be highly abundant and the size limit is set above the size at maturity. The main impact on Sand Flathead stocks is from the recreational sector with catches >20 times that of the commercial sector. A Sand Flathead monitoring program commenced in the summer of 2010/11; however, the time series is too brief at present to develop meaningful trends in relative abundance and there is a lack of historical data. These data are currently being analysed in anticipation of inclusion in the 2014/15 Scalefish assessment.

Increased interest in Sand Flathead by commercial operators is possible, and will likely be driven by rising market prices and/or reduced access and availability of other scalefish species. Future catch trends should be monitored closely along with those taken by the recreational sector. Given the possibility that Danish seine effort may increase it would be prudent to consider spatial management options to prevent regional concentration of effort on Sand flathead.

10. Flounder

Pleuronectidae family

STOCK STATUS	NOT ASSESSED
Greenback Flounder (<i>Rhombosolea tapirina</i>) constitute the majority of the commercial catches.	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends



Rhombosolea tapirina
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source												
Habitat	<ul style="list-style-type: none"> Sheltered sand. Between 0 and 100 m depth. 	Edgar (2008)												
Distribution	<ul style="list-style-type: none"> Southern Western Australia to southern New South Wales, and around Tasmania. Also in New Zealand. 	Edgar (2008)												
Diet	<ul style="list-style-type: none"> Polychaetes and benthic crustaceans. 	Edgar (2008) Ferguson (2006)												
Movement and stock structure	<ul style="list-style-type: none"> Several genetically distinct stocks in Australia: west coast of Tasmania, east coast of Tasmania, Victoria. 	van den Enden (2000)												
Natural mortality	<ul style="list-style-type: none"> Estimated at $M= 0.85$ (for New Zealand stock) 	Sutton et al. (2010)												
Maximum age	<ul style="list-style-type: none"> 10 years 	Sutton et al. (2010)												
Growth	<ul style="list-style-type: none"> Maximum length: 45 cm Maximum weight: 0.6 kg Growth (New Zealand) described by von Bertalanffy growth function $L = L_{\infty}(1 - e^{-k(t-t_0)})$ where L is the fork length (cm), t is the age (years), L_{∞} is the average maximum length for the species, k is a constant and t_0 is the (theoretical) age where length equals zero. Parameters estimates are : <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Sex</th> <th>L_{∞}</th> <th>k</th> <th>$t_{0\infty}$</th> </tr> </thead> <tbody> <tr> <td>Females</td> <td>55.82</td> <td>0.26</td> <td>-1.06</td> </tr> <tr> <td>Males</td> <td>52.21</td> <td>0.24</td> <td>-1.32</td> </tr> </tbody> </table> Length-weight relationship was estimated at $W = 0.036 L^{2.7}$ for females and $W = 0.039 L^{2.64}$ for males 	Sex	L_{∞}	k	$t_{0\infty}$	Females	55.82	0.26	-1.06	Males	52.21	0.24	-1.32	Edgar (2008) Kailola et al. (1993) Sutton et al. (2010)
Sex	L_{∞}	k	$t_{0\infty}$											
Females	55.82	0.26	-1.06											
Males	52.21	0.24	-1.32											

	where W is weight (g) and L is the tail length (cm).	
Maturity	<ul style="list-style-type: none"> Sexual maturity at about 218.6 mm TL for females and 190 mm TL for males. 	Crawford (1984)
Spawning	<ul style="list-style-type: none"> From June to October. Females are serial spawners and move from the shallows in deeper areas of tidal rivers and estuaries, and offshore for spawning. The relationship between batch fecundity and fork length is linear between 24.7 and 34.3 cm with $F = -1053.65 + 85.85L$, where F is the fecundity (in number of eggs) and L is the tail length (cm). Pelagic eggs, 0.7-1.0 mm in diameter 	Crawford (1984)
Early life history	<ul style="list-style-type: none"> Incubation of 82-93 hours Larvae hatch at 1.9 mm between May and November. Larvae remain in the plankton for over 30 days until they reach 6 mm, and then undergo metamorphosis which finishes 65 days post-hatching. Settlement inshore occurs during late winter to early summer. Juveniles live on sand flats in water less than a meter deep. 	Edgar (2008) Crawford (1984, 1986) Jenkins (1986)
Recruitment	<ul style="list-style-type: none"> No-stock recruitment relationship established. 	
Gillnet post release survival	<ul style="list-style-type: none"> High: 96.1% 	Lyle et al. (2014a)

Background

While the various Flounder species are not always differentiated in the logbooks, Greenback Flounder constitute the vast majority of the commercial catch. Long-snouted Flounder (*Ammotretis rostratus*) are also taken in small quantities. The main fishing methods used to target flounder are spear and gillnet. However, the ban in 2010 of setting unattended gillnets at night has essentially dissipated gillnet fishing for Flounder.

FISHING METHODS	Spear, graball net, some beach seine
MANAGEMENT METHODS	<p>Input control:</p> <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence) Recreational gear licence (graball and/or mullet net licence) <p>Output control:</p> <ul style="list-style-type: none"> Possession limit of 30 individuals for recreational Minimum size: 25 cm
MAIN MARKET	Local (Tasmania)

Current assessment

Catch, effort and CPUE

State wide Flounder landings have declined steadily since the mid-1990s, reaching 2.2 t in 2013/14 (Fig. 10.1A). Since the night gillnetting ban, Flounder are caught almost exclusively using spear. Commercial catches have contracted spatially over recent years and are now exclusively restricted to Norfolk Bay and the Tamar estuary (Fig. 10.2). Consistent with the trend in catches, effort for both methods has been steadily declining since the mid-1990's, with graball effort currently at minimal levels.

Catch rates based on gear units have remained relatively steady over time, although based on daily catches there has been a general decline for gillnets (Fig. 10.1C). There was, however, a minor upturn in catch rates for both methods during 2013/14.

Flounder is an important recreational species, and catches for the recreational sector exceed those of the commercial sector (Fig. 10.1A). Recreational catches were estimated at 15.2 t in 2000/01 (Lyle 2005), 10.1 t in 2007/08 (Lyle et al. 2009) and 7.2 t in 2012/13 (Lyle et al. 2014b). The majority of recreational fishing effort is centred in south-east Tasmania, in particular Norfolk-Frederick Henry Bay and D'Entrecasteaux Channel, as well as off northern Tasmania, especially the Tamar and north east coast regions.

Flounder





Figure 10.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. SP= spear, GN= graball net, NEC= north-east coast, SEC= south-east coast, EC= east coast.

A) Catch



B) Effort




Figure 10.2 (A) Flounder catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (44 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (27.1 t)	Yes	24.9 t (92.3%)
	• Catch increases by > 30% from previous year (>2.6 t)	No	
	• Catch decreases by > 30% from previous year (<1.4 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (1392 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0126 t/days fished)	Yes	0.0037 t/days fished (29.4%)

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (29.4 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (12.3 t)	Yes	10.2 t (82.6%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period (15.2 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (56.3% in 2007/08)	Yes	Latest estimate (2012/13): 77.4%
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0017)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status


NOT ASSESSED

The decrease in catches for the species is likely to be related to market demand but has also been impacted by the ban on overnight gillnetting. The Tasmanian catch is sold locally and demand for Flounder has decreased over the last two decades to the extent that the catch is so small now that its unreliable estimate of abundance.

11. Gould's Squid

Nototodarus gouldi

STOCK STATUS	SUSTAINABLE
High inter-annual variability in abundance. Commonwealth-managed species in Bass Strait (See: http://www.afma.gov.au/portfolio-item/goulds-squid/). Dual-endorsed vessels fish in Tasmanian waters especially in years of peak abundance.	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scalefish Fishery/Southern Squid-jig Fishery (Commonwealth)
INDICATOR(S)	Catch, effort and CPUE trends



Nototodarus gouldi
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> Open water. Inhabits coastal, inner shelf and shelf break waters down to 600 m depth. 	Stark (2008)
Distribution	<ul style="list-style-type: none"> From southern Queensland to mid-Western Australia, and around Tasmania 	Dunning (1998) Dunning and Förch (1998)
Diet	<ul style="list-style-type: none"> Small planktonic crustaceans, fish and squids 	Machida (1983) O'Sullivan and Cullen (1983) Smith (1983) Uozumi (1998)
Movement and stock structure	<ul style="list-style-type: none"> Move widely within a 300km² area but no large-scale migration between feeding and spawning areas as for other similar squid species Stock structure uncertain but appears to be only one population in southern Australia 	Jackson et al. (2005a) Triantafillos et al. (2004)
Natural mortality	<ul style="list-style-type: none"> No estimates available 	
Maximum age	<ul style="list-style-type: none"> Around 1 year 	Jackson et al. (2005b)
Growth	<ul style="list-style-type: none"> Maximum length: 40 cm. Maximum weight: 1.6 kg Size-at-age highly variable between individuals, years and locations. 	Norman and Reid (2000) Jackson et al. (2005b)

	<ul style="list-style-type: none"> Growth rate rapid: between 2.559 and 5.596 g.d⁻¹ for females, and between 1.622 and 5.307 g.d⁻¹ for males. 	Jackson et al. (2003)
Maturity	<ul style="list-style-type: none"> Size-at-50% maturity: between 30.6 to 31.4 cm ML for females, and 20.5 to 21.5 cm ML for males. 	Stark (2008)
Spawning	<ul style="list-style-type: none"> Spawns once and then die Spawning all year-round Egg mass are free-floating gelatinous sphere of at least 1.5 m in diameter and contains several thousands of eggs 	Jackson et al. (2005b) Uozumi (1998) O'Shea et al. (2004)
Early life history	<ul style="list-style-type: none"> Hatching throughout the year. 	Jackson et al. (2005b) Uozumi (1998)
Recruitment	<ul style="list-style-type: none"> Highly variable. No-stock recruitment relationship established. 	

Background

Gould's Squid, like most cephalopod species, has a very brief life cycle and can vary significantly in abundance between years. Environmental conditions are acknowledged as influences on larval and juvenile survival (Flood et al. 2012). It is likely that there is only one biological stock throughout southern Australian waters.

The Commonwealth Southern Squid-jig Fishery operates in Bass Strait waters using automatic squid-jigs and Gould's Squid are a regular by-product in the South East Trawl Fishery. Occasionally, Gould's Squid become available in high numbers in Tasmanian State waters, particularly around south-east Tasmania. Consequently, dual endorsed vessels may fish in State waters during the summer before moving back to traditional fishing grounds in Bass Strait.

Gould's Squid are processed into "tubes" and frozen. Given the unpredictable occurrence of the species in Tasmanian waters there is limited local processing capacity, which has limited the development of the fishery. There is also limited market demand with consumers preferring Southern Calamari. Due to the abovementioned factors, catch does not necessarily reflect biomass.

FISHING METHODS	Automated squid-jig, squid jig
MANAGEMENT METHODS	<p>Input control:</p> <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence, Automated squid-jig licence) Temporal and spatial closures (October-November) of some south east coast waters <p>Output control:</p> <ul style="list-style-type: none"> Possession limit of 15 individuals for recreational fishers
MAIN MARKET	Interstate

Current assessment

Catch, effort and CPUE

Gould's Squid availability in Tasmanian waters is highly variable, which is reflected in the catch history (Fig. 11.1A). Since 1995/96, there have been a few peaks of abundance, notably in 1999/2000, 2011/12 and 2012/13. The Gould's Squid catch for 2012/13 was the highest since 1995/96 (~1000 t) with the Australia-wide catch predominantly coming from Tasmanian waters

(Flood et al. 2014). By contrast in 2013/14 no Gould's Squid were landed by automatic jig in Tasmanian waters. Historically, in between the peaks, catches of the species have fluctuated between 0.1 and 50 t. The majority of the catch is taken in the south-east and east coast of Tasmania (Fig.11.2).

Gould's Squid catches from the recreational sector (Fig. 11.1A) were estimated at 5 t in 2000/01 (Lyle 2005), 36.6 t in 2007/08 (Lyle et al. 2009) and 21.4 t in 2012/13 (Lyle et al. 2014b).

Effort tends to follow the catch pattern closely and matches the availability of the species. In recent years high catches have been achieved with relatively low effort and a record catch due to high availability was obtained in 2012/13 (Fig. 11.1B).

Overall, catch rates remained relatively low until 2008/09. Since then, there has been an increase in catch rate, both in gear units and by days fished (Fig. 11.1C).

Gould's squid




Figure 11.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. AJ= automatic squid-jig, NEC= north-east coast, SEC= south-east coast, EC= east coast. Note: no notable catch and effort using Automatic squid-jig was recorded for 2005/06, 2006/07 or 2013/14.




Figure 11.2 (A) Gould's Squid catches (t) and (B) effort (days) by fishing blocks averaged from 2007/08 to 2011/12 (left) and during 2012/13 (right). No notable catch of Gould's Squid was recorded in 2013/14.

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (35.1 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (5.7 t)	Yes	5.7 t (100%)
	• Catch increases by > 30% from previous year (>1393.3)	No	
	• Catch decreases by > 30% from previous year (750.2 t)	Yes	750.2 t (100%)
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>238 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0048 t/days fished)	Yes	0.0023 t /days fished (47.4%)

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (39.6 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (2.1 t)	Yes	2.1 t (100%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period (5 t)	Yes	16.4 t (328%)
	• Proportion of recreational catch to total catch > previous proportion estimate (44.4% in 2007/08)	No	
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0656)	Yes	0.3 (-457%)
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

SUSTAINABLE

Many reference points were breached during the 2013/14 season as a result of the zero catch and lack of automatic squid-jig effort occurring in state waters. The lack of any effort in Tasmanian state waters suggests that the focus of the fishery was in Commonwealth waters. Spatial variability in abundance does not necessarily indicate a decrease in abundance in the southern Australian Gould's Squid stock.


A recreational reference point was also breached, however it is unlikely that the recreational sector poses a threat to the stock as estimated catches remain small in comparison to the commercial sector when Gould's Squid are available in Tasmanian waters and even more so when the Southern Squid-jig Fishery catches are taken into consideration.

Gould's Squid are short lived, spawn year round and display highly variable growth and size/age at maturity, meaning they can rapidly increase in numbers during favourable environmental conditions. As a result, Gould's Squid may be less susceptible to overfishing than longer lived species (Flood et al. 2012). However, their short life span (1 year) implies a reliance on a single cohort, which leaves the species very susceptible to environmental impacts on subsequent recruitment. The fishing effort in the Commonwealth Southern Squid-jig Fishery has decreased markedly since the late 1990's due to economic factors and catches are considered lower than the historical catches from the late 1970's. A biological stock depletion study found that no overfishing had occurred (Sahlqvist and Skirtun 2011). The species was classified as not overfished or subject to overfishing by the Fishery Status Reports 2012 (Woodhams et al. 2013), and as Sustainable by the Key Australian Fish Stock Reports (Flood et al. 2012, Flood et al. 2014). This assessment has therefore been applied to the Tasmanian component of the fishery.

12. Jack Mackerel

Trachurus declivis

STOCK STATUS	SUSTAINABLE
Shared stock with Commonwealth managed fishery (See: http://www.afma.gov.au/portfolio-item/jack-mackerel/).	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scafish Fishery/Small Pelagic Fishery (Commonwealth)
INDICATOR(S)	Catch, effort and CPUE trends



Trachurus declivis
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source								
Habitat	<ul style="list-style-type: none"> Open water. Between 0-500 m depth. 	Edgar (2008)								
Distribution	<ul style="list-style-type: none"> Western Australia (Shark Bay) to southern Queensland and around Tasmania. Also in New Zealand. 	Edgar (2008)								
Diet	<ul style="list-style-type: none"> Krill, planktonic crustaceans, fish 	Kailola et al. (1993)								
Movement and stock structure	<ul style="list-style-type: none"> Schooling fish. Normally live in continental shelf waters. May move close to seabed during winter. Most likely 2 subpopulations, one eastern Australian (east Tasmania and along the eastern seaboard of Australia) and one western Australian (west Tasmania, Great Australian Bight and Western Australia). 	Kailola et al. (1993) Bulman et al. (2008)								
Natural mortality	<ul style="list-style-type: none"> Estimated between $M = 0.63$ and 0.70 	Stevens and Hansfeld (1982)								
Maximum age	<ul style="list-style-type: none"> 25 years 	Paul (2000)								
Growth	<ul style="list-style-type: none"> Maximum length: 64 cm No difference between male and female growth. Growth described by von Bertalanffy growth function $L = L_{\infty}(1 - e^{-k(t-t_0)})$ where L is the length (cm), t is the age (years), L_{∞} is the average maximum length for the species, k is a constant and t_0 is the (theoretical) age where length equals zero. Parameters estimates are: <table border="1" data-bbox="395 1892 906 1966"> <thead> <tr> <th>Sex</th> <th>L_{∞}</th> <th>k</th> <th>$t_{0\infty}$</th> </tr> </thead> <tbody> <tr> <td>Combined</td> <td>36.2</td> <td>.267</td> <td>-1.21</td> </tr> </tbody> </table>	Sex	L_{∞}	k	$t_{0\infty}$	Combined	36.2	.267	-1.21	Paul (2000) Lyle et al. (2000)
Sex	L_{∞}	k	$t_{0\infty}$							
Combined	36.2	.267	-1.21							

	<ul style="list-style-type: none"> Length-weight relationship was estimated at $W = 1.46 \cdot 10^{-8} L^{2.982}$ for both males and females, where W is weight (g) and L is the length (cm). 	
Maturity	<ul style="list-style-type: none"> Sexual maturity between 3 and 4 years of age, at sizes around 27 cm and weights around 250 g. 	Webb (1976)
Spawning	<ul style="list-style-type: none"> Occurs over a wide area in Tasmania. Between late December and early March Pelagic eggs 	Stevens and Hansfeld (1982)
Early life history	<ul style="list-style-type: none"> Larvae carried by inshore currents Juveniles inhabit coastal and estuarine waters although they may sometimes be found offshore 	Kailola et al. (1993) Williams and Pullen (1986)
Recruitment	<ul style="list-style-type: none"> No-stock recruitment relationship established. 	
Gillnet post release survival	<ul style="list-style-type: none"> NA 	

Background

The Jack Mackerel fishery in Tasmania started in the early 1970's, with a one year venture catching 6 300 t in 1973. In 1985 another venture aimed at fishmeal production using purse seine nets and Jack Mackerel landings rose rapidly to reach 40 000 t in 1986/87 (Kailola et al. 1993). By 2000, fishers were struggling to catch surface schools and the industry began mid-water trawling for the species (and red bait) in Commonwealth waters. Small quantities of Jack Mackerel are also taken inshore as by-product of beach seining and inshore purse seining.

FISHING METHODS	Mainly purse seine, also beach seine
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence) Species licence (Mackerel A or B) Recreational gear licence (beach seine) Output control: <ul style="list-style-type: none"> Possession limit of 30 individuals for recreational Commercial catches taken by Mackerel licence holders (A & B) are decremented against the TAC allocated to the Commonwealth Small Pelagic Fishery
MAIN MARKET	Local (Tasmania)

Current assessment

Catch, effort and CPUE

Catches of Jack Mackerel in Tasmanian waters and reported in the catch and effort logbooks have been variable since 1995/96, oscillating between 2.6 and 59.8 t up until 2007/08 when there was a sharp increase in purse seine effort targeting Jack Mackerel (Fig. 12.1A). The Jack Mackerel catches peaked at 919.6 t in 2008/09, but declined sharply in 2010/11 and 2011/12 to around 60 t, the result of the major purse seiner ceasing operations. Catches usually focus on the south-east coast (Fig. 12.2). Only 0.4 t was recorded in Tasmanian waters in 2013/14. It should be noted that between 1995 and 1999, purse seine catches taken as part of the then Zone A Jack Mackerel fishery, and recorded in a separate logbook, ranged from 447 (1995/96)

to 8 458 t (1997/98), averaging 4 485 t per year for that period. These data are not presented in Fig. 12.1A.

Jack Mackerel is not a significant recreational species, catches estimated at 3.2 t in 2000/01 (Lyle 2005), 1.0 t in 2007/08 (Lyle et al. 2009) and 5.2 t in 2012/13 (Fig. 12.1A).

The use of purse seining by the major operator between 2008/09 to 2009/10 resulted in a spike in effort for that period of time. Beach seine effort has been slowly declining over time though Jack Mackerel represent a by-product and no meaningful catch rate trends can be drawn from these data (Fig. 12.1B). Purse seine catch rates were low until the species began being targeted in 2008/09 and remained high until 2011/12 which was the last year that the species was targeted (Fig. 12.1C). Since, landings have been low and there has been no targeted fishing in Tasmanian waters and few landings in the Commonwealth Small Pelagic Fishery (Flood et al. 2014).

Jack mackerel




Figure 12.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. BS= beach seine, PS= purse seine, SEC= south-east coast, EC= east coast. No catch for Jack Mackerel was reported during 2013/14.




Figure 12.2 (A) Jack Mackerel catches (t) and (B) effort (days) by fishing blocks averaged from 2007/08 to 2011/12. No notable catch for Jack Mackerel was reported during 2012/13 or 2013/14.

Limit reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (48.4 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (6.1 t)	Yes	5.7 t (93.4%)
	• Catch increases by > 30% from previous year (>0.21 t)	Yes	0.2 t (95.8%)
	• Catch decreases by > 30% from previous year (<0.11 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>239 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0191 t/days fished)	Yes	0.0082 t/days fished (43%)

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (26.2 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (8.6 t)	Yes	8.2 t (95.4%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period (3.2 t)	Yes	2.0 t (62.5%)
	• Proportion of recreational catch to total catch > previous proportion estimate (0.4% in 2007/08)	Yes	Latest estimate (2012/13: 96.3%)
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0254)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

SUSTAINABLE


The reference point for lowest catch, catch rate and value of the fishery were breached since very little fishing occurred for the species in 2013/14. The minimal commercial catch also meant that the recreational catch was proportionally higher than historically and therefore breached the recreational reference points. Recent trends in the commercial fishery have been the response to one operator entering and leaving the fishery and do not reflect the stock status. In fact a 2014 study assessed the spawning stock biomass for eastern Australia to be in the order of 150,000 tonnes (Ward et al. 2015). Jack Mackerel are assessed by the Commonwealth Small Pelagic Fishery Resource Assessment Group and based on current catch levels both the eastern and western Jack Mackerel stocks are assessed as not overfished and not subject to

overfishing (Woodhams et al. 2013, Flood et al. 2014) and this assessment has been applied to the Tasmanian component of the fishery.

13. Jackass Morwong

Nemadactylus macropterus

STOCK STATUS	SUSTAINABLE
Shared stock with Commonwealth-managed fishery (See: http://www.afma.gov.au/portfolio-item/jackass-morwong/)	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scalefish Fishery/Southern and Eastern Scalefish and Shark Fishery (Commonwealth)
INDICATOR(S)	Catch, effort and CPUE trends



Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> Exposed sand and silt, reefs. Between 5 and 400 m depth. 	Edgar (2008)
Distribution	<ul style="list-style-type: none"> From mid Queensland to southern Western Australia, and around Tasmania. Also in New Zealand, southern Africa and South America. 	Edgar (2008)
Diet	<ul style="list-style-type: none"> Polychaete worms, crustaceans, molluscs and echinoderms 	Godfriaux (1974)
Movement and stock structure	<ul style="list-style-type: none"> No genetic variation in southern Australia indicating larval mixing. Existence of at least 3 sub-populations: Tasmania, New South Wales/Victoria and Great Australian Bight 	Richardson (1982) Grewe et al. (1994) Elliott and Ward (1994) Thresher et al. (1994)
Natural mortality	<ul style="list-style-type: none"> $M = 0.10$ (New Zealand population) 	Parker and Fu (2011)
Maximum age	<ul style="list-style-type: none"> 50 years 	Edgar (2008)

Growth	<ul style="list-style-type: none"> Maximum length: 70 cm TL Maximum weight: 2.9 kg Growth varies according to location. Growth described by von Bertalanffy growth function $L = L_{\infty}(1 - e^{-k(t-t_0)})$ where L is the length (cm FL), t is the age (years), L_{∞} is the average maximum length for the species, k is a constant and t_0 is the (theoretical) age where length equals zero. Parameters estimates are: <table border="1"> <thead> <tr> <th>Sex</th> <th>L_{∞}</th> <th>k</th> <th>t_0</th> </tr> </thead> <tbody> <tr> <td>Females</td> <td>38.4</td> <td>0.36</td> <td>-0.07</td> </tr> <tr> <td>Mal s</td> <td>36.2</td> <td>0.42</td> <td>0.15</td> </tr> </tbody> </table>	Sex	L_{∞}	k	t_0	Females	38.4	0.36	-0.07	Mal s	36.2	0.42	0.15	Francis (2001) Kailola et al. (1993) Jordan (2001a)
Sex	L_{∞}	k	t_0											
Females	38.4	0.36	-0.07											
Mal s	36.2	0.42	0.15											
Maturity	<ul style="list-style-type: none"> Sexual maturity at about 25 cm TL and about 3 years of age 	Edgar (2008)												
Spawning	<ul style="list-style-type: none"> Between February and June At least 2 spawning areas: northern one (probably southern New South Wales and eastern Victoria) and a southern one (probably western and southern Tasmania). 	Lyle and Ford (1993) Bruce et al. (2001a)												
Early life history	<ul style="list-style-type: none"> Planktonic larval stage of 7-10 months. Larvae up to 30 mm drift with current on the surface up to 250 km east of Tasmania Settlement at 7-9cm long. Juveniles live near shallow reefs 	Francis (2001) Bruce et al. (2001a) Kailola et al. (1993)												
Recruitment	<ul style="list-style-type: none"> No-stock recruitment relationship established. 													
Gillnet post release survival	<ul style="list-style-type: none"> Moderate: 52% 	Lyle et al (2014a)												

Background

Jackass Morwong is a Commonwealth-managed species. While there is a good market for Jackass Morwong, the species is not available in large numbers in Tasmanian waters and is hence a by-product of graball netting rather than a target species. Tasmanian commercial catches reached a maximum of around 250 t in the late 1980's due to the inshore trawl fishery, but this fishery has since ceased operations and State catches have declined significantly. Most of the Jackass Morwong catch originates from trawling outside Tasmanian waters and while the stocks were assessed to be overfished from 2008 to 2010, in 2012 the stock was classified as not overfished nor subject to overfishing (Woodhams et al. 2013). The species was not assessed in the 2014 Status of Australian Fish Stocks.

FISHING METHODS	Mainly graball net, also hand line and drop-line
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence) Recreational licence (graball and/or mullet net) Output control: <ul style="list-style-type: none"> Possession limit of 15 individuals for recreational Minimum size: 25 cm TL
MAIN MARKET	Local

Current assessment

Catch, effort and CPUE

Total commercial catch of Jackass Morwong was 1 t in 2013/14 (Fig 13.1A). Commercially, Jackass Morwong is mainly caught by gillnet and landings have declined steadily since 1995/96. The majority of the catch is taken from the east and south-east coast (Fig. 13.2).

Jackass Morwong is an important recreational species, with catches estimated at higher levels than those of the commercial fishery (Fig. 13.1A). Estimates were 31.9 t in 2000/01 (Lyle 2005), 6.8 t in 2007/08 (Lyle et al. 2009), 7.7 t in 2011/12 (Tracey et al. 2013) and 16.1 t in 2012/13 (Lyle et al. 2014b).

A decline in effort of both gear units and days fished has mirrored the declining catches for the species (Fig. 13.1B). Catch rates have slowly declined since the early 2000's but stabilised since 2007/08 at a low level (Fig. 13.1C).

Jackass morwong





Figure 13.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. GN= graball net, SEC= south-east coast, EC= east coast.

A) Catch



B) Effort




Figure 13.2 (A) Jackass Morwong catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (136.9 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (19 t)	Yes	17.9 t (94.2%)
	• Catch increases by > 30% from previous year (>2.2 t)	No	
	• Catch decreases by > 30% from previous year (<1.2 t)	Yes	0.6 t (35%)
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>1057 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0095 t/days fished)	Yes	0.0051 t/days fished (53.6%)

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (19 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (13.1 t)	Yes	12.1 t (92.2%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period (31.9 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (64.2% in 2007/08)	Yes	Latest estimate (2012/13): 88.5%
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period • (-0.0017)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

SUSTAINABLE

The lowest catch reference point was breached, which reflects the ongoing declining commercial catches in Tasmanian waters. Linked to this decline, one of the recreational indicators was breached, indicating that the proportion of recreational catch to total catch had increased in 2012/13 since the last recreational survey (2007/08).

A single east Australian stock of Jackass Morwong is shared between the Commonwealth and Tasmania with catch and catch rates declining in Commonwealth fisheries in a fashion similar to that observed in the Tasmanian fishery. This was driven by a prolonged period of reduced recruitment that is believed to be a result of climate induced changes to ocean currents in eastern Tasmania (Wayte 2013). Due to the extended larval phase of Jackass Morwong, these changes have meant that larvae are often dispersed away from Tasmanian waters with successful recruitment being correlated with years where oceanographic conditions are

comparable to historical averages (Wayte 2013). While the Jackass Morwong stocks were considered overfished in the late 2000s, the latest Fishery Status Reports in 2011 and 2012 reclassified the stocks as not overfished and not subject to overfishing (Woodhams et al. 2013, Flood et al. 2014). This assessment was due to a reduction of catches for the species as a response to management action in the Commonwealth fishery and revision of the stock assessment model. The total catch (recreational and commercial) of Jackass Morwong for Tasmania is low compared to the Commonwealth catch and the stocks are most likely shared. Although there has not been a formal Commonwealth assessment since 2012, unpublished Commonwealth statistics indicate the east coast stock is rebuilding under current total allowable catch levels and there has not been further reductions required. As the Fishery Status Reports describe the stock biomass as sustainable and fishing mortality as sustainable, we choose to accept their ranking given the Tasmanian fishery shares the same stocks.

14. Leatherjacket

Monacanthidae family

STOCK STATUS	UNDEFINED
Various species of Leatherjackets are found inshore around Tasmania's coastline and are not differentiated in logbooks. Leatherjackets are a by-product species and are not actively targeted. There is little biological information available for most species.	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends



Leatherjacket
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> Seagrass and reefs. Down to 200 m depending on species. 	Edgar (2008)
Distribution	<ul style="list-style-type: none"> Australia is the centre of diversity for this family with more than half of the estimated 90 species occurring here, mainly in temperate areas. 	Edgar (2008)
Diet	<ul style="list-style-type: none"> Epiphytes attached to seagrass, algae, fish flesh, molluscs and crustaceans depending on species. Many species are omnivorous. 	FishBase (2013)
Movement and stock structure	<ul style="list-style-type: none"> Many species are site-attached 	Barrett (1995b)
Natural mortality	<ul style="list-style-type: none"> Undefined for most species 	
Maximum age	<ul style="list-style-type: none"> No information 	
Growth	<ul style="list-style-type: none"> Maximum length: from 90 mm to 600 mm 	Edgar (2008)
Maturity	<ul style="list-style-type: none"> Little information 	
Spawning	<ul style="list-style-type: none"> Little information 	
Early life history	<ul style="list-style-type: none"> Little information 	
Gillnet post release survival	<ul style="list-style-type: none"> High: 95% 	Lyle et al (2014a)

Background

Leatherjackets are a by-product of fish traps and they are also caught in netting operations but are generally discarded. While Leatherjackets are consumed on the mainland, there is little market demand for them in Tasmania.

FISHING METHODS	Mainly fish trap, also graball net and handline
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none">• Gear licence (Scalefish fishing licence)• Recreational gear licence (graball and/all mullet net) Output control: <ul style="list-style-type: none">• Possession limit of 15 individuals for recreational fishers• Minimum size: 20 cm
MAIN MARKET	Local

Current assessment

Catch, effort and CPUE

Leatherjacket catches have declined continuously since the early 2000's, reaching a minimum of 2.4 t in 2012/13 before increasing slightly to 2.9 t in 2013/14 (Fig. 14.1A). While catches from graball nets have consistently remained at low levels since 1995/96, catches from fish traps have declined over time. Leatherjackets are caught on the east, north-east and north-west and south-east coasts (Fig. 14.2).

Leatherjackets are also caught by the recreational sector, with catch estimates in recent surveys at a similar level to commercial catches (Fig. 14.1A). Estimates were 8.2 t in 2000/01 (Lyle 2005), 2.6 t in 2007/08 (Lyle et al. 2009), 2.3 t in 2009/10 and 1.8 t in 2012/13 (Lyle et al. 2014b).

Both fish trap and graball fishing effort has decreased through time (Fig. 14.1B). Fish trap effort (in gear units and fishing day) dropped significantly around 2007/08. Abalone gut was the preferred bait used in fish traps; however a ban on its use has been in place since 2008 to prevent the spread of an abalone virus (AVG). Following this ban, fishing effort using fish traps declined as other baits are considered inferior.

Catch rates in gear unit and days fished have remained relatively stable over time for both graball nets and fish traps and are presently at levels similar to those at the beginning of the time series (Fig. 14.1C).

Leatherjacket





Figure 14.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. FP= fish trap, GN= graball net, SEC= south-east coast, EC= east coast, NEC= north-east coast.

A) Catch



B) Effort




Figure 14.2 (A) Leatherjacket catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (27.7 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (12.2 t)	Yes	9.3 t (76.2%)
	• Catch increases by > 30% from previous year (>3.1 t)	No	
	• Catch decreases by > 30% from previous year (<1.6 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>1607 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (0.0046 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (16.5 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (10.4 t)	Yes	7.5 t (72.4%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period (8.2 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (38.0% in 2007/08)	Yes	Latest estimate (2012/13): 41.9%
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0014)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

UNDEFINED

The very low landings in 2013/14 meant that the lowest catch, lowest fishing mortality and the proportion of recreational to commercial catch reference points were breached. These decreased catches are, however, the result of a decline in the use of fish traps and a lack of demand.


Leatherjackets tend to be site-attached and have limited home ranges. Two decades of monitoring eastern Tasmanian Marine Protected Areas (MPA) show that there is no significant difference in Leatherjacket abundances inside or outside of MPA's (e.g. brown striped Leatherjacket, Toothbrush Leatherjacket). Leatherjackets also have a very high post release survival following capture in gillnets (Lyle et al. 2014a). These results, along with the low landings, suggest that fishing is unlikely to have a significant impact on Leatherjacket

populations. There is however too little information at the species level to confidently classify the status of Leatherjacket stocks.

15. Longsnout Boarfish

Pentaceropsis recurvirostris

STOCK STATUS	NOT ASSESSED
Little information is available for this species.	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends



Pentaceropsis recurvirostris
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> Exposed reef. Between 4 and 260 m depth. 	Edgar (2008)
Distribution	<ul style="list-style-type: none"> From mid New South Wales to southern Western Australia, and around Tasmania. 	Edgar (2008)
Diet	<ul style="list-style-type: none"> Brittle stars, polychaetes and brown algae 	Edgar (2008) Scott et al. (1974)
Movement and stock structure	<ul style="list-style-type: none"> Unknown 	
Natural mortality	<ul style="list-style-type: none"> Unknown 	
Maximum age	<ul style="list-style-type: none"> Unknown 	
Growth	<ul style="list-style-type: none"> Maximum length: 61 cm 	Edgar (2008)
Maturity	<ul style="list-style-type: none"> Unknown 	
Spawning	<ul style="list-style-type: none"> Unknown 	
Early life history	<ul style="list-style-type: none"> Unknown 	
Gillnet post release survival	<ul style="list-style-type: none"> High: 99.7% 	Lyle et al (2014a)

Background

Boarfish are a by-product of graball netting. Due to the large minimum legal size and the requirement to release undersized fish, boarfish are also a regularly discarded. Boarfish are considered good eating and there is a small local market for them. Shark nets also catch boarfish as a by-product in State waters; these catches have been reported to the Commonwealth since 2000/01.

FISHING METHODS	Graball net, shark net in the past
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none">• Gear licence (Scalefish fishing licence)• Recreational gear licence (graball and/or mullet net) Output control: <ul style="list-style-type: none">• Trip limit of 50 kg for commercial fishers• Possession limit of 2 individuals for recreational fishers• Minimum size: 45 cm
MAIN MARKET	Mainly local

Current assessment

Catch, effort and CPUE

In Tasmania, boarfish catches are now exclusively derived from graball net (Fig. 15.1A). Catches have been declining through time but appear to have stabilised since 2008/09 with landings in 2013/14 of 1 t, which is lower than the previous year (2.1 t) (Fig. 15.1A). Catches are taken exclusively from the east and south-east coasts (Fig. 15.2). Boarfish are not caught by rod and line and no recreational catch estimates are available for the graball and spearfishing catch of this species. However, fewer than 1000 individuals were recorded (both kept and released) in the 2012/13 survey (Lyle et al. 2014b) indicating that boarfish are not a common recreational species.

After peaking in 1997/98, commercial graball net effort has remained stable (Fig. 15.1B). Catch rates have remained relatively stable over time (Fig. 15.1C).

Boarfish





Figure 15.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. SN= shark net, GN= graball net, SEC= south-east coast, EC= east coast.

A) Catch



B) Effort




Figure 15.2 (A) Boarfish catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (10.4 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (7.2 t)	Yes	6.2 t (86.1%)
	• Catch increases by > 30% from previous year (>2.7 t)	No	
	• Catch decreases by > 30% from previous year (< 1.5 t)	Yes	0.4 t (27.4%)
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>858 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0038 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (6.2 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (3.6 t)	Yes	2.6 t (72.3%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period	Not estimated	
	• Proportion of recreational catch to total catch > previous proportion estimate	Not estimated	
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0009)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status


NOT ASSESSED

Reference points associated with low catches were breached. Catches are lower now due to a lesser quantity of gillnetting effort than occurred in the reference period. There is too little information available to assign a stock status for this species. Boarfish is a non-target species that is taken in very small quantities. In addition to catches taken in State waters, there is also a by-product fishery from Commonwealth shark netting activity. The high size limit means that the majority of individuals captured are released and the species has very high post-release survival (Lyle et al. 2014a).

16. Yelloweye Mullet

Aldrichetta forsteri

STOCK STATUS	Not Assessed
Commercial and recreational catches are low and at similar levels.	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends



Aldrichetta forsteri
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> Sheltered sand, seagrass, up to 20 m depth. May ascend rivers into freshwaters. 	Edgar (2008)
Distribution	<ul style="list-style-type: none"> Western Australia (Shark Bay) to New South Wales, and around Tasmania, New Zealand. 	Edgar (2008)
Diet	<ul style="list-style-type: none"> Planktonic animals for juveniles, benthic crustaceans and molluscs for medium-sized fish and almost exclusively algae for larger fish 	Edgar (2008)
Movement and stock structure	<ul style="list-style-type: none"> Schooling fish No genetic studies but there appears to be 2 populations (eastern Australia and western Australia) 	Kailola et al. (1993)
Natural mortality	<ul style="list-style-type: none"> M estimated at 0.66 (New Zealand) 	Paul and Taylor (1998)
Maximum age	<ul style="list-style-type: none"> 7 years 	Curtis and Shima (2005)
Growth	<ul style="list-style-type: none"> Maximum length of 50 cm. Maximum weight: 950 g Differential growth between males and females, and between locations Growth described by von Bertalanffy growth function $L = L_{\infty}(1 - e^{-k(t-t_0)})$ where L is the fork length (cm), t is the age (years), L_{∞} is the average maximum length for the species, k is a constant and t_0 is the (theoretical) age where length equals zero. 	Edgar (2008) Curtis and Shima (2005) Gorman (1962) Last et al. (1983) Chubb et al. (1981)

	Parameters estimates are:									
	<table border="1"> <thead> <tr> <th>Sex</th> <th>L_{∞}</th> <th>k</th> <th>$t_{0\infty}$</th> </tr> </thead> <tbody> <tr> <td>Combined</td> <td>40</td> <td>0.51</td> <td>-0.03</td> </tr> </tbody> </table>	Sex	L_{∞}	k	$t_{0\infty}$	Combined	40	0.51	-0.03	
Sex	L_{∞}	k	$t_{0\infty}$							
Combined	40	0.51	-0.03							
	<ul style="list-style-type: none"> Length-weight relationship was estimated at $W = 0.000239 L^{3.2}$ for females and males combined where W is weight (g) and L is the fork length (cm). 									
Maturity	<ul style="list-style-type: none"> 2-3 years 	Kailola et al. (1993)								
Spawning	<ul style="list-style-type: none"> Form large aggregation prior to spawning Spawn in coastal waters in summer and autumn, probably in estuaries Fecundity between 125,000 and 630,000 eggs Pelagic eggs 	Chubb et al. (1981) Kailola et al. (1993)								
Early life history	<ul style="list-style-type: none"> Juveniles enter estuaries and sheltered bays when they are 3-4 cm long, and remain there until they reach 25-30 cm tail length. As they grow older, animals gradually move to more open coastal waters 	Kailola et al. (1993)								
Gillnet post release survival	<ul style="list-style-type: none"> Low: 10% 	Lyle et al (2014a)								

Background

Mullet are occasionally targeted by netting, specifically beach and purse seining as well as small mesh net. The vast majority of catch is Yelloweye Mullet, but it is possible that some of the catch includes Sea Mullet (*Mugil cephalus*). Mullet are also targeted by recreational fishers using rod and line or small mesh gillnets called 'mullet nets'.

FISHING METHODS	Mostly beach seine, also small mesh net (mullet net for recreational) and purse seine
MANAGEMENT METHODS	<p>Input control:</p> <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence, small mesh gillnet licence) Recreational gear licence (graball and/or mullet net) <p>Output control:</p> <ul style="list-style-type: none"> Possession limit of 15 individuals of each species for recreational Minimum size: 25 cm
MAIN MARKET	Mostly local

Current assessment

Catch, effort and CPUE

Mullet are caught using beach seine and small mesh nets. After peaking in 1999/2000, Yelloweye Mullet catches have decreased and stabilised at less than 5 t since 2007/08. The catch for 2012/13 was 9.2 t, which was similar to the landings in the early 2000's and was due to an increase in beach seine catch in the east and south-east coasts. The catch in 2013/14 catches dropped to 2.9 t, a level comparable to that for most other recent years (Fig. 16.1A). Recreational catches were estimated at 6.5 t in 1996/97, 30 t in 2000/01, 1.7 t in 2009/10 and 7.1 t in 2012/13 (Lyle et al. 2014b)(Fig. 16.1A).

Beach seine effort was relatively stable until 2005/06 after which it declined rapidly and remained at low levels (Fig. 16.1B). Purse seine effort has been more variable with no clear pattern with the trends dictated by the targeting of Australian Salmon.

Catch rates for beach seining have remained relatively constant over time, whereas catch rates for purse seining are highly variable with no apparent pattern, probably because this species is rarely targeted (Fig. 16.1C).

Mullet





Figure 16.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. BS= beach seine, PS= purse seine, EC= east coast, SEC= south-east coast and NEC= north-east coast.

A) Catch



B) Effort




Figure 16.2 (A) Yelloweye Mullet catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (31.2 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (10.8 t)	Yes	7.9 t (73.1%)
	• Catch increases by > 30% from previous year (>12.0 t)	No	
	• Catch decreases by > 30% from previous year (<6.4 t)	Yes	3.5 t (54.3%)
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>327.8 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0107 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (14.5 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (7.3 t)	Yes	4.4 t (60.1%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period (30.0 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (73.3% in 2007/08)	No	
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0061)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

NOT ASSESSED

Other than in 2012/13, catches of Yelloweye Mullet have been stable at low levels for the past five years, following a decrease in effort in the traditional fishing grounds in northern Tasmania. The high catch in 2012/13 meant that several reference points were breached in 2013/14 when catches had declined back to levels similar to those for 2007/08 to 2011/12. Despite the low commercial and recreational catches, there is too little information available to confidently classify the status of Yelloweye Mullet in Tasmanian waters.

17. Pike

Longfin Pike *Dinolestes lewini*

Shortfin Pike *Sphyraena novaehollandiae*

STOCK STATUS	NOT ASSESSED
The main species targeted is the Shortfin Pike, also known as Snoek or Snook. Little information is available on either of the species.	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends

Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> <u>Longfin Pike</u>: exposed reef. Down to depth of 60 m. <u>Shortfin Pike</u>: exposed reef, sand, seagrass, offshore waters. Down to depth of 20 m. 	Edgar (2008)
Distribution	<ul style="list-style-type: none"> <u>Longfin Pike</u>: Western Australia to New South Wales, and around Tasmania <u>Shortfin Pike</u>: Western Australia to southern Queensland, and northern Tasmania. 	Edgar (2008)
Diet	<ul style="list-style-type: none"> <u>Both species</u>: fish 	Coleman and Mobley (1984) Scott et al. (1974)
Movement and stock structure	<ul style="list-style-type: none"> <u>Longfin Pike</u>: no information <u>Shortfin Pike</u>: highly migratory pelagic species that often occurs in shoals of 50 or more individuals 	Kailola et al. (1993)
Natural mortality	<ul style="list-style-type: none"> <u>Both species</u>: no information 	
Maximum age	<ul style="list-style-type: none"> <u>Both species</u>: no information 	
Growth	<ul style="list-style-type: none"> <u>Longfin Pike</u>: Maximum length of 90 cm. <u>Shortfin Pike</u>: Maximum length of 1.1 m, maximum weight of 5.6 kg, maximum age of about 20 years. 	Edgar (2008) Kailola et al. (1993)
Maturity	<ul style="list-style-type: none"> <u>Longfin Pike</u>: no information <u>Shortfin Pike</u>: 42 cm length 	Bertoni (1995)
Spawning	<ul style="list-style-type: none"> <u>Longfin Pike</u>: no information <u>Shortfin Pike</u>: assumed to take place from October to January 	Kailola et al. (1993)
Early life history	<ul style="list-style-type: none"> No information 	
Gillnet post release survival	<ul style="list-style-type: none"> NA 	

Background

While commonly referred to as Pike, Longfin Pike (*Dinolestes lewini*) and Shortfin Pike (*Sphyræna novaehollandiæ*) are not related and come from two separate families. Pikes are mainly targeted by troll and small-mesh net (north coast only), and are also a by-product of beach seining and graball netting. While there is a local and interstate market for Shortfin Pike, Longfin Pike are of lesser demand. There are some uncertainties about the correct reporting of the two species in logbooks. The vast majority of Pike catches are likely to be Shortfin Pike, which is confirmed by anecdotal reports from the industry.

FISHING METHODS	Troll, also beach seine, graball net and small mesh net
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none">• Gear licence (Scalefish fishing licence)• Recreational gear licence (graball and/or mullet net) Output control: <ul style="list-style-type: none">• Possession limit of 15 individuals of each species for recreational
MAIN MARKET	Local and interstate (Victoria)

Current assessment

Catch, effort and CPUE

Both species were pooled together because of uncertainties in the logbook reporting. Pike catches were variable but followed a relatively stable trend from 1995/96 to the mid-2000s. However, catches declined over the last six years to reach a record low of 8.2 t in 2012/13 (Fig. 17.1A). In 2013/14 catches were 9.9 t and somewhat similar to other years in the recent past. Although there was a peak in Pike catches in the south-east in 1996/97 and 1997/98, catches traditionally originate from the north-west coast (Fig. 17.2). There are no estimates of recreational landings (by weight) but evidence suggests that Pike are not a major recreational target (Lyle et al. 2009, Lyle and Tracey 2012) and that around 57% of all Pike caught by recreational fishers are released (Lyle et al. 2009). Nevertheless, in 2012/13, 3,895 Pike were landed by recreational fishers (Lyle et al. 2014b) and if an average weight of 1 kg per fish is assumed, the landings in 2012/13 would have been at least 4 t.

After increasing until 2001, troll effort for Pike has decreased and stabilised (Fig. 17.1B). Beach seine effort in both gear unit and days fished has remained stable over time but Pike represent a by-product of this fishing method rather than a target.

Catch rates in gear units have remained stable for both gear types, whereas catch rates in days fished has increased over time for trolling (Fig. 17.1C).

Pike





Figure 17.1 A) Annual commercial catch (t) by gear (left) and region (right). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. BS = beach seine, TR = troll, MN = small mesh gillnet, NWC = north-west coast.

A) Catch



B) Effort




Figure 17.2 (A) Pike catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (18.8 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (9.5 t)	No	
	• Catch increases by > 30% from previous year (>10.7 t)	No	
	• Catch decreases by > 30% from previous year (<5.8 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>519.2 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0104 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (18.3 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (13.3 t)	Yes	3.4 t (25.5%)
	• Latest recreational catch estimate > recreational catch estimate from the reference period (based on numbers)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate	Not estimated	
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0024)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

NOT ASSESSED

The lowest catch reference point was breached, reflecting the low troll and beach seine catches for the current assessment year. The fishery for Pike is relatively small and mainly limited to the northern part of Tasmania. Moreover, the species is rarely targeted recreationally. Due to stable catch rates and the above factors, Pike are unlikely to be overfished; however, as they represent a by-product, or target in mixed species fisheries, catch rate may be unreliable and there is consequently too little information available to confidently estimate the status of the stock.

18. School Whiting

Sillago flindersi

STOCK STATUS	SUSTAINABLE
This species was previously commonly referred to as the Southern School Whiting (<i>Sillago bassensis</i> , also known as Western School Whiting) but the species present in Tasmania and caught commercially is in fact the Eastern School Whiting (<i>Sillago flindersi</i>). Commonwealth managed species (See: http://www.afma.gov.au/portfolio-item/eastern-school-whiting/)	
IMPORTANCE	Minor
STOCK(S)	Tasmanian Scalefish Fishery/Southern and Eastern Scalefish and Shark Fishery (Commonwealth)
INDICATOR(S)	Catch, effort and CPUE trends



Species biology

Parameters	Estimates	Source								
Habitat	<ul style="list-style-type: none"> Costal lakes, estuaries and along outer coast. Down to 170 m depth. 	Gomon et al. (2008)								
Distribution	<ul style="list-style-type: none"> Endemic to south-eastern Australia, from southern Queensland to western Victoria, and around Tasmania 	Gomon et al. (2008)								
Diet	<ul style="list-style-type: none"> Feed mainly on crustaceans, amphipods, decapods, mysids and copepods. Juveniles consume mostly copepods. 	Burchmore et al. (1988)								
Movement and stock structure	<ul style="list-style-type: none"> There is some evidence of 4 genetically distinct stocks (2 in New South Wales, 1 in Tasmania and 1 in Victoria). Commonwealth assessments state that the evidence for separate stocks is weak and manage the species as single stocks. 	Dixon (1987) Morison et al. (2012)								
Natural mortality	<ul style="list-style-type: none"> No information but likely to be around $M = 0.7$ based on related species. 	Butcher and Hagedoorn (2003)								
Maximum age	<ul style="list-style-type: none"> 7 years 	Kailola et al. (1993)								
Growth	<ul style="list-style-type: none"> Maximum length: 33 cm SL Growth described by von Bertalanffy growth function $L = L_{\infty}(1 - e^{-k(t-t_0)})$ where L is the length (cm), t is the age (years), L_{∞} is the average maximum length for the species, k is a constant and t_0 is the (theoretical) age where length equals zero. Parameters estimates are: <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Sex</th> <th>L_{∞}</th> <th>k</th> <th>t_0</th> </tr> </thead> <tbody> <tr> <td>Combined</td> <td>23.9</td> <td>0.46</td> <td>-0.50</td> </tr> </tbody> </table>	Sex	L_{∞}	k	t_0	Combined	23.9	0.46	-0.50	Gomon et al. (2008) Tilzey (1994)
Sex	L_{∞}	k	t_0							
Combined	23.9	0.46	-0.50							

Maturity	<ul style="list-style-type: none"> Reached at 2 years and a size of 14-16 cm FL 	Hobday and Wankowski (1987) Burchmore et al. (1988)
Spawning	<ul style="list-style-type: none"> Spring to late summer. Females release between 30 000 and 110 000 eggs in total during the season 	Hobday and Wankowski (1987)
Early life history	<ul style="list-style-type: none"> Juveniles inhabit inshore waters. 	FishBase (2013)
Gillnet post release survival	<ul style="list-style-type: none"> NA 	

Background

There is a large Commonwealth fishery for School Whiting that has landed 1000–2500 t for the last 30 years with about 75% of the catch from Commonwealth waters being taken by Danish seine operating out of Lakes Entrance, Victoria (Morison et al. 2012). The New South Wales state managed fishery has been increasing in recent years and now accounts for around 60% of the total catch, which has led to some equity disagreements as their catch is deducted from the Commonwealth total allowable catch (Morison et al. 2012). School Whiting have been exploited in Tasmania since the mid-1970's with catches ranging from 20–175 t throughout the 1980's (Kailola et al. 1993). The vast majority of the catch is taken by Danish seine in the south of the State. Danish seine fishing operations target either School Whiting (with Flathead as a by-product) or Flathead (with School Whiting as by-product), which accounts for the often opposing trends in catches for the two species. School Whiting are mainly marketed and processed in Melbourne.

FISHING METHODS	Danish seine
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence, small mesh gillnet) Danish seine licence (with whiting codend endorsement) Output control: <ul style="list-style-type: none"> Possession limit of 30 individuals for recreational fishers
MAIN MARKET	Interstate

Current assessment

Catch, effort and CPUE

Catches have been variable but relatively stable over time, with a yearly average of 35.1 t between 1995/96 and 2013/14 (Fig. 18.1A). Landings were at their lowest in 2011/12 and 2012/13 (16 – 17 t) but increased to 37.2 t in 2013/14. Catches were concentrated on the south-east coast as has historically been the case (Fig. 18.2). Recreational catches are low compared to commercial catches, and were estimated at 0.8 t in 2000/01 (Lyle 2005), 3.4 t in 2007/08 (Lyle et al. 2009) and 2.1 t in 2012/13 (Lyle et al. 2014b) (Fig. 18.1A).

Effort in both gear unit and days fished has been decreasing steadily over time, though there was an increase in 2013/14 (Fig. 18.1B). Catch rates by gear units and days fished have generally increased since 1995/96, though there are fluctuations (Fig. 18.1C), which are likely caused by changes in target species (i.e. School Whiting versus Tiger Flathead).

Whiting





Figure 18.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. DS= Danish seine, SEC= south-east coast.

A) Catch



B) Effort




Figure 18.2 (A) School Whiting catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (152.3 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (25.5 t)	No	
	• Catch increased by > 30% from previous year (>21.4 t)	Yes	15.7 t (73.2%)
	• Catch decreased by > 30% from previous year (>11.5 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>210.1 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0145 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (42.7 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (30.6 t)	No	
	• Latest recreational catch estimate > recreational catch estimate from the reference period (0.8 t)	Yes	1.3 t (162%)
	• Proportion of recreational catch to total catch > previous proportion estimate (8.7% in 2007/08)	No	
Biomass	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0664)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

SUSTAINABLE

The >30% increase in catch from previous year reference point was breached, however, this may not be significant in a commercial fishery that is dependent on the interstate market and is (generally) based on a single operator, who does not work full time. These patterns are also dictated by the species being targeted (Tiger Flathead or School Whiting).

While the most recent recreational catch was higher than during the reference period, catches by the recreational sector remain low and are inconsequential at the scale of the stock.


The most recent Commonwealth stock assessment was carried out using data up until 2010 (see stock assessment summary by Morison et al. (2012)). While outdated, this stock assessment showed increasing catch rates and given a total allowable catch is in place, which has not been taken in recent years due to a decline in exports because of the high Australian dollar it is unlikely that the species has been recruitment overfished in recent years. Based, on

this evidence, and catch rate trends from New South Wales, Victoria (and Tasmania), the most recent Status of Australian Fish Stocks ranked School Whiting as sustainably fished (Flood et al. 2014) and we followed this assessment.

19. Southern Calamari

Sepioteuthis australis

STOCK STATUS	UNDEFINED
Vulnerability of calamari to fishing pressure is unclear but probably high. High recreational interest in the species.	
IMPORTANCE	Key
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch and CPUE trends



Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> Shallow inshore water. 	Gomon et al. (2008)
Distribution	<ul style="list-style-type: none"> Endemic to southern Australia and northern New Zealand waters 	Gomon et al. (2008)
Diet	<ul style="list-style-type: none"> Various crustaceans and fishes 	Norman (2000)
Movement and stock structure	<ul style="list-style-type: none"> Highly mobile. Undergoes migration between feeding grounds and spawning grounds. Preliminary genetic studies showed there are a minimum of 5 genetically distinct stocks in Australia. 98% of the Tasmania population belongs to a single genetic stock which is also found in various proportions in the South Australian, New South Wales and Western Australian populations, suggesting some degree of interbreeding between these areas. A recent, and more thorough, genetic study showed a single stock across southern Australia with Tasmania being particularly important in terms of reproduction. 	Triantafillos & Adams (2001) Triantafillos (2004) Smith <i>et al.</i> (2015)
Natural mortality	<ul style="list-style-type: none"> High. Embryo mortality rate between 5% and 25%. 	Steer et al. (2004)
Maximum age	<ul style="list-style-type: none"> Short-lived (<1 year). Maximum recorded ages are 275 days for males and 263 days for females. 	Pecl et al.(2004)
Growth	<ul style="list-style-type: none"> Rapid growth at 7-8% body weight/day in individuals <100days old, decreasing to 4-5% BW/day in squids older than 200 days. No gender difference in growth. Growth from 80 days (when recruitment to inshore waters occurs) best described by a power function $L = 2e^{-6t^{3.5332}}$ where L is the mantle 	Pecl et al.(2004) Triantafillos (2004) Data from Pecl (2004)

	<p>length (mm) and t is the age (days).</p> <ul style="list-style-type: none"> Extremely variable growth. Some variability may be explained by temperature and food availability (individuals hatched in warmer season/year generally grow faster) but there is also a genetic component. Length-weight relationship was set at $W = 0.00081L^{2.427}$ where W is weight (g) and L is the dorsal mantle length (mm). 	
Maturity	<ul style="list-style-type: none"> Size-at-50% maturity estimated at 184.5mm for females. 	Data from Pecl (2006)
Spawning	<ul style="list-style-type: none"> Major spawning period in spring/summer (March to July) in Tasmania, with low levels of spawning occurring all year round Great Oyster Bay (east coast Tasmania) a known spawning ground. Spawning aggregations are male-biased. Multiple spawners with individual spawning activity occurring over several months (up to 3.5 months). Females deposit eggs together in collective egg masses, attaching the finger-like capsules to the substrate by small stalks. 	<p>Moltschaniwskyj & Pecl (2003) Pecl et al. (2004)</p> <p>Pecl et al. (2006)</p>
Early life history	<ul style="list-style-type: none"> Incubation time estimated at 4 to 8 weeks depending on water temperature. Hatchlings (2.4-7 mm) swim to the surface and can be found near spawning grounds for 20-30 days. Habitat and ecology between 20-80 days unknown. From 80-150 days, juveniles are found in deeper water adjacent to spawning ground. Individuals become available to the fishery between 90-120 days of age. 	<p>Steer et al. (2002)</p> <p>Pecl (2000)</p> <p>Pecl (2004)</p>

Background

The fishery for Southern Calamari initially developed in the mid-1990s in Great Oyster Bay and then expanded rapidly to the south (including Mercury Passage, Maria Island and Tasman Peninsula) during the latter half of the 1990s, with annual catches rising from less than 20 t prior to 1995/96 to around 90 t in 1998/99. Since then, catches have fluctuated between 40 and 110 t. The expansion of the fishery was accompanied by a massive increase in effort, particularly squid-jig, which has become the primary capture method in recent years. Calamari are taken in lesser quantities by purse seine, beach seine, spear and dipnet. Although some night fishing occurs, Calamari are mainly targeted during the day over shallow areas of seagrass and macro-algae where they aggregate to spawn.

FISHING METHODS	Squid-jig (main), purse seine, beach seine, spear, dipnet
MANAGEMENT METHODS	<p>Input control:</p> <ul style="list-style-type: none"> Gear licence (Scalefish fishing licence) Species licence (Southern Calamari licence) for the South-east region Danish seine licence Class seine licences Temporal and spatial closures (mid-October to mid-November) of some south east coast waters

MAIN MARKET	Output control:
	Trip limit of 10 calamari in 24 hour period for SE waters for scalefish licence holders not the holder/operator of a calamari or seine licence Local and interstate

Current assessment

Catch, effort and CPUE

The total commercial catch of Southern Calamari for 2013/14 was 67.5 t (Fig. 19.1A). While the main fishery is concentrated off the central east and south-east coasts, catches are reported from all areas apart from the west coast (Fig. 19.2). Spawning area closures first introduced in 1999 appear to have been successful in reducing pressure on the main spawning grounds (Great Oyster Bay and Mercury Passage) and in encouraging industry to spread their effort throughout the State (Fig. 19.1A). Catches increased off the north-east coast (around Bicheno and Waterhouse Island) in 2002/03 (Fig. 19.1A) and off the north-west coast (mainly around Stanley) in 2007/08, with current production levels similar to that for the Mercury Passage and the south-east coast (Fig. 19.2).

Effort is focused in Great Oyster Bay, Mercury Passage on the east coast, around the Tasman Peninsula in the south-east, and Stanley and Ringarooma Bay in the north (Fig. 19.2). There has been a general shift in effort in recent years from Great Oyster Bay to Mercury Passage (Fig. 19.2) where catch rates have been more consistent over time (Fig. 19.1D). Low availability of Calamari in Great Oyster Bay has been confirmed by industry reports for several years now. These observations are consistent with surveys of egg production from the main inshore spawning beds in Great Oyster Bay that indicated very low egg densities for 2011 and 2012, and continues the general decline in egg production since 2004 (Fig. 19.3). Assuming egg production is related to the abundance of adults on spawning grounds, then the general decline in catch rates in Great Oyster Bay (Fig. 19.1D) is likely to be influenced by both the impacts of closures during peak fishing periods and by a reduced spawning stock size. Industry has however reported large number of egg masses outside the Great Oyster Bay area, e.g. around Maria Island and in the north of the State, suggesting that there are other spawning grounds providing recruitment for the stock. Despite the above, CPUE remains relatively stable in the nearby Mercury Passage and has, if anything, increased slightly in recent years (19.1D).

Effort and CPUE have been relatively stable for the north-east coast since its initial increase in 2002/03, while effort and CPUE continued to increase until 2008/09 before stabilising in recent years for the north-west coast (Fig. 19.1C and E).

The most recent recreational catch estimate (2012/13) indicated that landings were 63.5 t (Lyle et al. 2014b) and similar to commercial landings. This is a considerable increase from previous years with landings from the recreational fishery increasing consistently throughout the last 13 years (Fig. 19.1A) suggesting increased interest in the species.














Fig. 19.1 A) Annual commercial catch (t) by gear (left) and by region (right), and best estimates of recreational catches (single squares); Commercial squid-jig effort based on B) gear units (left) and days fished (right) relative to 1998/99 for MP and 1997/98 for GOB and C) gear units (left) and days fished (right) relative to 2001/02 for NEC and NWC; Commercial squid-jig catch per unit effort (CPUE) based on D) weight per gear unit (left) and weight per day (right) relative to 1998/99 for MP and relative to 1997/98 for GOB, and E) weight per gear unit (left) and weight per day (right) relative to 2001/02 for NEC and NWC; SJ = squid-jig, DN = dipnet; NWC= north-west coast, NEC= north-east coast, SEC = south-east coast, MP = Mercury Passage, GOB = Great Oyster Bay. Only years with >5 operators are shown.

A) Catch



B) Effort




Figure 19.2 (A) Calamari catches (t) and (B) effort (days) by fishing blocks averaged from 2007/08 to 2012/13 (left) and during 2013/14 (right). South East Waters management boundaries are also displayed.




Figure 19.3 Calamari egg production estimated from dive surveys of the main inshore spawning beds in Great Oyster Bay by week of year. No survey was performed in 2013 or 2014.

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > highest catch from the 1990/91 to 1997/98 range (33 t)	Yes	34.5 t (104.5%)
	• Catch < lowest catch from the 1990/91 to 1997/98 range (5.8 t)	No	
	• Catch increases by > 30% from previous year (>83.1 t)	No	
	• Catch decreases by > 30% from previous year (<44.8 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>879 days fished)	Yes	294.1 days fished (36.8%)
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0114 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (104.8 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (33 t)	No	
	• Catch variation from the previous year above the greatest inter-annual increase from the reference period (67.9 t)	No	
	• Catch variation from the previous year above the greatest inter-annual decrease from the reference period (-69.7 t)	No	
	• Latest recreational catch estimate > recreational catch estimate from the reference period (17.7 t)	Yes	47.6 t (269%)
	• Proportion of recreational catch to total catch > previous proportion estimate (30.3% in 2007/08)	Yes	Latest estimate (2012/13): 51.3%
Biomass	• CPUE < 3 rd lowest CPUE value from the reference period (0.0198 t/days fished)	No	
	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0030)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

UNDEFINED

The current reference range for catch and effort are based on a period prior to the development of the fishery, and therefore result in a triggering of the reference points. The proposed

reference points overcome this shortfall by using a more appropriate reference range. None of the proposed indicators were breached for commercial catches. The recreational reference points were breached as a result of continued increasing interest in the species from this sector, with the recreational catch of similar magnitude to that of the commercial sector, indicating that overall fishing pressure on the species has increased over the past decade or so, despite relative stability in commercial landings.

Vulnerability of Calamari to fishing pressure is unclear but probably high because spawning aggregations are targeted and the species has an annual or sub-annual life span that renders the stock susceptible to spawning and/or recruitment failure. The species-specific licence has effectively capped commercial effort in the south east and, if the population is allowed to spawn (during the fishing closures) prior to the main harvest period, the population may be able to sustain high rates of fishing mortality without detrimental effects on future recruitment in this area. There remains high, and increasing, interest in the species from the recreational sector, particularly in areas in south-eastern waters such as D'Entrecasteaux Channel (closed to commercial fishing), Norfolk and Frederick Henry Bay, and Great Oyster Bay. While catch rates have slightly improved in Great Oyster Bay following a strong decline in catches and catch rates in 2008/09 and 2009/10, there is a need to monitor the developments in the fishery over the coming years, especially in light of the decreasing egg production in one of the main known spawning areas. The identification of source and sink populations supporting the Tasmanian fishery is critical to ensure the sustainable use of this resource. Particularly because a recent genetic study showed that Tasmania was the most important state for providing recruitment throughout southern Australia (Smith et al. (2015)), which may show the positive influence the spawning closures have had as they are not in place in other states.

Catch and effort have increased over time outside of the South East Waters (defined as waters between Whale Head to Lemon Rock for calamari management), particularly in the central east and north-west coasts. Virtually nothing is known about the status of the stocks off the north coast but growing interest from commercial and recreational fishers suggests a need for at least some baseline monitoring. Stable or increasing catch rates suggest that current levels of fishing are not excessive but the status will need to be monitored in future, particularly because anecdotal evidence suggests increasing recreational fishing effort in the region as well.

Due to the many uncertainties listed above, there is currently too little information available to confidently classify the status of this stock.

20. Southern Garfish

Hyporhamphus melanochir

STOCK STATUS	Sustainable
After a strong decline in catches from 2006/07 coupled with changes in population structure, which prompted management actions, the species is showing early signs of recovery.	
IMPORTANCE	Key
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends; changes in size/age composition



Hyporhamphus melanochir
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> Shallow inshore water (<20m depth) in association with seagrass beds. 	Gomon et al. (2008)
Distribution	<ul style="list-style-type: none"> Eden (New South Wales) to Perth (Western Australia), including Bass Strait and Tasmanian waters. Endemic to Australia. 	Gomon et al. (2008)
Diet	<ul style="list-style-type: none"> Predominantly herbivores (seagrass, algal filaments). Also consume planktonic crustaceans, worms, diatoms and stray insects landing on the surface. 	Edgar (2008), Klumpp and Nichols (1983)
Movement and stock structure	<ul style="list-style-type: none"> Schooling fish, highly mobile. School near the surface at night and close to bottom during day. There are 4 genetically distinct populations distributed in Western Australia, western South Australia, eastern South Australia/Victoria and Tasmania. It is likely that at least two Garfish subpopulations exist in Tasmania, as suggested by the different size and age characteristics exhibited through sampling of the north and east coasts. This has not been confirmed genetically. 	Grant (1991) St Hill (1996) Jones et al. (2002) Hartmann & Lyle (2011)
Natural mortality	<ul style="list-style-type: none"> High. Estimated at 55% for adults of four years and over for the east coast population. 	Jones (1990)
Maximum age	<ul style="list-style-type: none"> Up to 9 years old. In recent years, individuals 6 years and older have been rare in catches. 	Jordan et al. (1998) Hartmann & Lyle (2011)
Growth	<ul style="list-style-type: none"> From 6 month onwards, growth follows a Von Bertalanffy growth function with $L_{\infty} = 34.3$, $k = -0.54$ and $t_0 = 0.23$ Length-weight relationship: $W = 0.0011L^{3.4403}$ where W is weight (g) and L is the fork length (cm). 	Jordan et al. (1998) Hartmann & Lyle (2011)
Maturity	<ul style="list-style-type: none"> Size-at-50% maturity estimated at 19.9cm for females and 17.1cm for males. The relationship between batch fecundity and fork length is linear with $F = 188.75L - 3585.8$, where F is the fecundity (in number of eggs) and L is the fork length (cm). 	Hartmann & Lyle (2011)
Spawning	<ul style="list-style-type: none"> Spawning is concentrated in shallow (<5 m deep) over 	Jordan et al. (1998)

	<p>beds of drift algae in eastern Tasmania. It occurs over an extended period of at least five months from October to February, with peak activity occurring between October and December.</p> <ul style="list-style-type: none"> Eggs are around 2.93 mm in diameter and are negatively buoyant, sinking to the bottom after fertilisation and becoming attached to drift algae. 	
Early life history	<ul style="list-style-type: none"> Incubation time estimated at one month. Large hatchlings (7.8-8.5mm) are assumed to stay in shallow waters as evidenced by the presence of small juvenile (0+ cohort) in the coastal sheltered waters of east Tasmania. 	Jordan et al. (1998)
Recruitment	<ul style="list-style-type: none"> Variable. No-stock recruitment relationship established. 	
Gillnet post release survival	<ul style="list-style-type: none"> NA 	

Background

Traditionally a winter beach seine fishery, catches were initially centred off the northeast coast, including Flinders Island. More recently the fishery has extended to the east and southeast coasts and with the introduction of dip-nets, catches have increasingly been taken over the summer months. Nowadays, Garfish is caught almost exclusively by beach seine on the north-east coast and mainly by dip-nets off the south-east and east coasts.

FISHING METHODS	Mainly dip-net and beach seine
MANAGEMENT METHODS	<p>Input control:</p> <ul style="list-style-type: none"> Temporal closure during spawning: mid-Nov to mid-Dec for southern waters, mid-Jan to mid-Feb for northern waters (Northern and southern waters are delineated by a line following the north coast of Tasmania, joined and bounded in the west by a line of latitude through Cape Grim and in the east by a line of latitude through Cape Naturaliste). Gear restriction (Scalefish fishing licence, Purse seine licence, Beach seine licence) <p>Output control:</p> <ul style="list-style-type: none"> Legal size: 25 cm (upper jaw to end of tail) Possession limit of 30 (recreational fishers)
MAIN MARKET	<ul style="list-style-type: none"> Mainly local

Current assessment

Biological characteristics

The sharp and unexpected decline in catches in 2006/07 and 2007/08 prompted the need to better understand stock status and resulted in the introduction of a one month closed season in 2009 to protect the species during the spawning season.

Size composition

Post the mid-1990s there was a reduction of 2–3 cm in the average size of Garfish in eastern Tasmania, and up to 5–6 cm in northern Tasmania (Fig. 20.3). Annual length frequency distributions tended to be unimodal with a peak between 28–33 cm in the 1990s and 27–28 cm during the late 2000s. These reductions were associated with a sharp decline in commercial catches and may be indicative of a more general change in population structure. Since 2011 however, there appears to be a recovery with average size in eastern Tasmania moving towards 30 cm.

A similar pattern was apparent in the north coast. During the mid-1990s individuals between 35 and 40 cm were relatively common in the catch with modal peaks >30 cm but in subsequent years fish larger than about 35 cm were rare and modal peaks were generally <30 cm (Fig. 20.3). Since 2012, there has been an increase in the average size of Garfish, and in the catch of larger fish (>35 cm).

Age composition

The age structure of east and north coast catches taken between 2008 and 2011 were similar, and both were dominated by 2–4 year old fish (Fig. 20.2). In both the east and the north coasts, these dominant classes accounted for between 80 and 90% of the catch numbers annually. Fish aged up to 7 years and 8 years were represented in catches from the north and east coast respectively, although individuals 6 years or older were rare (<2% numbers for the north and <8% for the east coast). By contrast, the age structure of east coast catches taken between 1995–1997 were characterised by a broader range of age classes, 3–7 year olds, with 4 and 5 year olds dominating and accounting for 43% of the total sample (Jordan et al. 1998). Over the past decade there was no evidence for changed gear selectivity (e.g. via gear modifications), nor have there been reported changes in fishing practices or market preference that would account for the shift to younger/smaller fish. The contraction in age structure and sharp fall in catches since the mid-2000s may therefore reflect fundamental changes in population structure, which may have resulted from heavy fishing pressure and/or poor recruitment during the early 2000s.

In 2012, however, the age distribution changed in both regions, with a lesser proportion of two year olds in the catch and fish >3 years old becoming more prevalent, indicating the start of a recovery (Fig. 20.2). The age distribution in the east coast is now dominated by 3–5 years old, which represented over 75% of the catch numbers. The age distribution in the north is still dominated by 2–4 years old (80% of the catch numbers) but in both regions there is an increase in the proportion of older fish with individuals 6 years or older accounting for around 13% in numbers.

Catch, effort and CPUE

The total commercial catch of Garfish for 2013/14 was 34.3 t, which is less than the previous five years catch (40.2–77.3 t) (Fig. 20.3A). After many years of relative stability in Garfish catches at 80–90 t per annum, catches fell sharply in 2006/07 and have fluctuated between 40–70 t since that time. Catches, both historically and in 2013/14, are concentrated off the north-east coast (Fig. 20.3A and Fig. 20.4A). Historically, there were also moderate landings in the east and south-east, however these were greatly reduced in 2013/14 (Fig. 20.3A and Fig. 20.4A).

Recreational Garfish catches are low compared to commercial catches and were estimated around 2 t in 2000/01, 2007/08 and 2012/13 (Henry and Lyle 2003, Lyle et al. 2009, Lyle et al. 2014b).

Effort of both commercial gear types has been slowly decreasing through time and there was a further notable reduction in 2013/14 (Fig. 20.3B and Fig. 20.4B). Dip-net effort initially increased to a peak during 1998/99 but has subsequently decreased to a lower level (Fig. 20.3B, days fished), while beach seine effort experienced a more recent decline and reached a historic minimum in 2013/14.




Figure 20.1. Size composition of Southern Garfish by year for east coast (left) and north coast (right) commercial catches. *n* is the sample size.




Figure 20.2. Age composition of Southern Garfish by year for east coast (left) and north coast (right) commercial catches.

Catch rates for beach seine fluctuated much more strongly over time than those for dip-net but catch rates for both beach seine and dip-net increased from 2011/12–2012/13 before declining slightly in 2013/14 (Fig. 20.3C). Catch rates remain at close to the historic maxima for both gears. Beach seine catch rates fluctuated at high levels until 2006/07, when they stabilised at low levels until 2010/11. By contrast, dip-net catch rates have undergone two periods of decline with a recovery period in between each. In the context of schooling species such as Garfish, catch rates may be relatively insensitive to changes in abundance.





Figure 20.3 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. BS= beach seine, DN= dip-net, SEC= south-east coast, EC= east coast, NEC= north-east coast.

A) Catch



B) Effort




Figure 20.4 (A) Garfish catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (91.6 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (56.2 t)	Yes	21.9 t (39%)
	• Catch increases by > 30% from previous year (>66.9 t)	No	
	• Catch decreases by > 30% from previous year (<26 t)	Yes	1.8 t (5%)
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>1081 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0306 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (91.6 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (66.2 t)	Yes	32 t (48.3%)
	• Catch variation from the previous year above the greatest inter-annual increase from the reference period (35.5 t)	No	
	• Catch variation from the previous year above the greatest inter-annual decrease from the reference period (-43.6 t)	No	
	• Latest recreational catch estimate > recreational catch estimate from the reference period (1.9 t)	Yes	0.1 t (5.3%)
	• Proportion of recreational catch to total catch > previous proportion estimate (3.7% in 2007/08)	Yes	Latest estimate (2012/13): 3.8%
Biomass	• CPUE < 3 rd lowest CPUE value from the reference period (0.05 t/days fished)	No	
	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0084)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	No	
	• Significant numbers of unhealthy fish landed	No	

Stock status

SUSTAINABLE

The low catch in 2013/14 breached several reference points (low catch and rapid decline from previous year) reflecting that catches are still low compared to levels prior to 2006/07. Catch rates, however, remain high and it appears to be a decline in effort by the beach seine sector,

in particular, that has resulted in the low 2013/14 landings. There has also been a notable reduction in the catch from the east and south-east of the state. While the recreational reference points were also breached, catches have remained essentially unchanged and are insignificant when compared to the commercial catch.

Vulnerability of Southern Garfish to fishing pressure is unclear but is probably moderate to high due to the schooling behaviour of the species, which means they can be efficiently targeted under favourable conditions. As a relatively short-lived species, and with a population dominated by relatively few age classes, any recruitment variability is likely to have a marked impact on population size.

There was some clear evidence of a reduction in average fish size and a truncation of the age structure up to 2011 that may be indicative of a more general change in population structure from heavy fishing pressure and/or poor recruitment. There were early signs of population recovery in 2012 (increasing size and age), which may be linked to the implementation of the spawning closure for commercial operators in 2009. The catch rates during the last two years imply some degree of stock recovery. Further, the current minimum legal length of 25 cm total length is above the size at maturity for both sexes and therefore likely to offer a relatively high degree of protection to the population (size at 50% maturity is 17.1 cm fork length for males and 19.9 cm for females, Hartmann and Lyle 2011).

Some industry members have expressed concern about the effects of dip-nets on the schooling behaviour of Garfish. Specifically, it has been suggested that intense dip-net activity tends to cause schools to break and hence reduce opportunities to use beach seines to target the species, possibly affecting catch rates. Since such interactions tend to be localised, analyses at the spatial resolution of fishing blocks are unlikely to be sensitive enough to detect such impacts.

The low commercial catch, historically high catch rates and an increase in the number of year classes represented in the catch implies suggests that the stock is not subject to heavy fishing pressure and as such, Southern Garfish are considered a sustainable stock.

21. Striped Trumpeter

Latris lineata

STOCK STATUS	TRANSITIONAL-RECOVERING
After a lack of recruitment for over a decade, Striped Trumpeter is showing signs of recovery in the last four years. However, the stock currently relies exclusively on recent recruits.	
IMPORTANCE	Key
STOCK(S)	Tasmanian Scalefish Fishery, Commonwealth fisheries
INDICATOR(S)	Catch, effort and CPUE trends



Latris lineata
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> Exposed reefs and rocky bottom down to 300m depth. 	Edgar et al. (2004) Gomon et al.(2008)
Distribution	<ul style="list-style-type: none"> Sydney (New South Wales) to Albany (Western Australia), Tasmania, New Zealand, Amsterdam Islands (southern Indian Ocean) and most of temperate Southern hemisphere (excl. South Africa and South America). 	Edgar et al. (2004) Gomon et al.(2008)
Diet	<ul style="list-style-type: none"> Small fish, cephalopods, crustaceans. 	Nichols et al. (1994)
Movement and stock structure	<ul style="list-style-type: none"> Juveniles have limited movement, remaining around shallow reefs for several years before moving into deeper offshore reefs. Adults have the capacity to undergo wide-scale movements (e.g. Tasmania to St Paul Island in Indian Ocean) Uniform stock structure in Tasmanian waters (no significant genetic separation of populations) 	Tracey and Lyle (2005) Lyle and Jordan (1999) Lyle and Murphy (2001) Tracey et al. (2007b)
Natural mortality	<ul style="list-style-type: none"> Estimated at $M=0.1$. 	Tracey and Lyle (2005)
Maximum age	<ul style="list-style-type: none"> Estimated at 43 years. 	Tracey and Lyle (2005)
Growth	<ul style="list-style-type: none"> Maximum length: 1.2 m Maximum weight: 25 kg Rapid juvenile growth (mean FL = 28 cm after 2 years, 42 cm after 4 years). Slower adult growth (large range of size- 	Gomon et al.(2008) Murphy and Lyle (1999)

	<p>at-age over 50 cm FL)</p> <ul style="list-style-type: none"> Growth for both sex described by a two-phase von Bertalanffy growth function $L = \left(1 - \int_{t=-t_0}^{+t^\delta} \frac{1}{\sigma\sqrt{2\pi}} e^{\left(\frac{-(t-t_{01})^2}{2\sigma^2}\right)} \right) (L_{\infty 1}(1 - e^{-k_1(t-t_{01})}) + \varepsilon) + \left(1 - \int_{t=t^\delta}^{+t_{max}} \frac{1}{\sigma\sqrt{2\pi}} e^{\left(\frac{-(t-t^\delta)^2}{2\sigma^2}\right)} \right) (L^\delta + (L_{\infty 2} - L^\delta)(1 - e^{-k_2(t-t_{02})}) + \varepsilon)$ <p>where L is the length (mm), t is the age (years), $L_{\infty 1}$ and $L_{\infty 2}$ are the average maximum length for the species for the 1st and 2nd growth phase respectively, k_1 and k_2 are constants, t_{01} and t_{02} are the (theoretical) age where length equals zero for the 1st and 2nd growth phase respectively, L^δ and t^δ are the length and age of transference from one growth phase to the next, t_{max} is the maximum age present in the sample, σ^2 is the standard deviation of cumulative density function with mean t^δ, and ε is an error term.</p> <p>Parameters estimates are:</p> <table border="1"> <thead> <tr> <th>$L_{\infty 1}$</th> <th>k_1</th> <th>t_{01}</th> <th>L^δ</th> <th>$L_{\infty 2}$</th> <th>k_2</th> <th>t_{02}</th> <th>t^δ</th> <th>σ^2</th> </tr> </thead> <tbody> <tr> <td>532.77</td> <td>0.43</td> <td>0.03</td> <td>450.1</td> <td>871.59</td> <td>0.08</td> <td>3.49</td> <td>4.4</td> <td>1.0</td> </tr> </tbody> </table> <ul style="list-style-type: none"> Length-weight relationship for both sex was estimated at $W = 2E^{-5}L^{3.00}$, where W is weight (g) and L is the fork length (mm). 	$L_{\infty 1}$	k_1	t_{01}	L^δ	$L_{\infty 2}$	k_2	t_{02}	t^δ	σ^2	532.77	0.43	0.03	450.1	871.59	0.08	3.49	4.4	1.0	Tracey and Lyle (2005)
$L_{\infty 1}$	k_1	t_{01}	L^δ	$L_{\infty 2}$	k_2	t_{02}	t^δ	σ^2												
532.77	0.43	0.03	450.1	871.59	0.08	3.49	4.4	1.0												
Maturity	<ul style="list-style-type: none"> Size-at-50% maturity estimated at 54 cm FL (62 cm TL; 6.8 years) for females and 53 cm FL (61 cm TL; 6.2 years) for males. Batch fecundity (BF) estimate: $BF = 4.15E^{-8}FL^{4.69}$, where FL is fork length in cm. 	Tracey et al. (2007a) IMAS unpublished data																		
Spawning	<ul style="list-style-type: none"> July to early October depending on geographic location (early start and finish at lower latitudes). Multiple spawners, highly fecund (100,000 to 400,000 eggs for females weighing 3.2 kg and 5.2 kg respectively). Small pelagic eggs (1.3 mm diameter). 	Ruwald et al. (1991) Ruwald, 1992 (1992) Hutchinson (1993)																		
Early life history	<ul style="list-style-type: none"> Complex and extended larval phase of at least 9 months before settlement. No information on size and timing of settlement. Juveniles of around 18 cm FL (23 cm TL) have been caught on shallow reefs off south-east coast in January. 	Ruwald et al. (1991) Ruwald, 1992 (1992) Murphy and Lyle (1999)																		
Recruitment	<ul style="list-style-type: none"> Highly variable. No-stock recruitment relationship established. 	Murphy and Lyle (1999)																		
Gillnet post release survival	<ul style="list-style-type: none"> NA 																			

Background

Striped Trumpeter have a long history of commercial exploitation in Tasmania, being highly valued for its eating qualities. There is also a high level of interest in the species from recreational fishers and charter boat operators. The species is taken by a variety of fishing gears, with hooks and gillnets being the primary methods. Juvenile Striped Trumpeters are taken predominantly by graball net in inshore waters and usually in depths <50 m, whereas adult fish are taken in deeper offshore waters by hook methods (drop-line, handline, bottom longline, trotline) and as by-product in large mesh gillnets (shark nets). Catches are concentrated off the east coast, including Flinders Island, as well as off the south and south-west coasts of Tasmania.

Responsibility for the management of Striped Trumpeter was passed to Tasmania in 1996 through an Offshore Constitutional Settlement (OCS) arrangement with the Commonwealth. A memorandum of understanding accompanied the OCS, specifying trip limits for Commonwealth only fishers. As part of the Tasmanian scalefish management plan, gear restrictions for all commercial scalefish fishers operating in State waters were introduced in 1998. This, however, enabled dual endorsed operators (i.e. holders of a Tasmanian licence and a Commonwealth permit for southern shark or South East Non Trawl fisheries) as well as rock lobster fishers to take unrestricted quantities of Striped Trumpeter in offshore waters using their gear allocations. In 2000, the State Government introduced a combined trip limit of 250 kg for Striped Trumpeter, yellowtail kingfish and snapper for all fishers (Commonwealth and State) in inshore and offshore waters relevant to Tasmania to limit the potential for expansion of effort directed at these species. Over time, there has been additional management measures targeted at the species, including a spawning closure, a decrease in the recreational possession limit and several increases in the minimum size limit for the species (currently 50 cm total length (TL), which is below the size at maturity of 62 cm TL for females and 61 cm TL for males).

Although the seasonal closure is not supported by the Commonwealth managed fisheries, in 2013 they reduced their Striped Trumpeter trip limit component to 150 kg year around (it is still a part of the 250 kg combined striped trumpeter/snapper/yellowtail kingfish bundle, but only a maximum of 150 kg can comprise striped trumpeter).

FISHING METHODS	Mainly handline, also graball net and drop-line
MANAGEMENT METHODS	<p>Input control:</p> <ul style="list-style-type: none"> • Gear licence (Scalefish fishing licence) • Temporal closure (Sept–Oct) <p>Output control:</p> <ul style="list-style-type: none"> • Trip limit of 250 kg • Trip limit of 150 kg for commonwealth operators • On water bag limit of 4 fish for recreational fishers • Possession limit of 8 fish for recreational fishers • Minimum size (500 mm TL)
MAIN MARKET	Mainly local

Current assessment

Biological characteristics

Size composition

In 1999, the size distribution was unimodal and dominated by 50 cm fork length (FL) fish, with a tail of older fish up to 89 cm length (Fig. 21.1). Many of the subsequent sample sizes were low and sampling was opportunistic, which may not represent population size structure. Years with

good sample sizes, however, suggest that no significant recruitment occurred for about a decade during the late 1990s and early 2000s following a major recruitment event in 1993. This 1993 cohort still dominated the catches in 2009 and at this time, samples were dominated by fish around 62 cm in length. In 2010, the size distribution changed and was again dominated by 50 cm fish, suggesting a new wave of recruits had entered the fishery (Fig. 21.1). The size structure appears stable since 2010 suggesting recruitment over at least 3 consecutive years. The fishery is currently relying heavily on these recent recruits. The lack of individuals over 60 cm in 2012 is almost certainly an artefact of sampling.

Age composition

The very strong 1993 year class (5 year olds in 1999, 6 year olds in 2000, etc.) was prominent in samples obtained from research fishing and commercial catches up until 2009 (Fig. 21.2). As for size, the low sample sizes in many years may not represent population age structure. Notwithstanding these concerns, the data demonstrated a consistent lack of recent recruits (3–5 year olds) up until 2010 which, along with catch declines, implied an extended period of low recruitment. During this time the population appears to have been sustained largely by strong year classes spawned during the 1990s.

Samples from 2010 to 2012 were dominated by 4–6 year olds (the age at which the species tends to recruit to the offshore hook fishery), providing evidence of recent recruitment success. While the relative strength of these cohorts is unknown, they provide a positive sign for the future of the stock, although caution is recommended since previous indicators have suggested that the adult segment of the population may still be in a depleted state (due to fishing and lack of recruitment over many years). No individuals older than 10 years old were detected in the 2012 samples but this reflects the low sample size rather than a major contraction in the population age structure.

Catch, effort and CPUE

The recent catch history in waters south of latitude 39° 12'S (i.e. waters incorporated within the OCS agreement for Striped Trumpeter), including catches reported in the Victorian and Commonwealth logbooks, is presented in Table 21.1. In the early 1990s catches by Victorian vessels were significant, peaking at around 37 t. Since the mid-1990s, data from this sector have been unavailable, though it is assumed that subsequent catches have been reported in Commonwealth logbooks. Other than 1999/00 when >14 t was taken, Commonwealth catches have been relatively low since that time.

Annual production was high at over 110 t in the early 1990's, Victorian vessels accounting for 17–39% of the total, but then fluctuated between 70–80 t through the mid-1990's before increasing again to over 100 t by the late 1990's (Table 21.1). Catches almost halved in 2000/01 to less than 50 t and have remained low since that time. This trend was observed by all methods in Tasmania (Fig. 21.3A). Despite increases in production between 2010/11 and 2011/12 the catch declined to 10.3 t in 2013/14, which is the lowest catch reported since records have been kept.

Commonwealth catches are believed to be substantially underreported and the lack of precise estimates of recreational catch represents sources of uncertainty in estimating the total mortality. The recreational fishery has heavily targeted Striped Trumpeter in the past, with an estimated 38 t in 2000/01 (Lyle 2005). Estimates of recreational catches in 2007/08 were highly uncertain but likely to be lower, with an estimated 19 t for combined catches of striped and Bastard Trumpeter (Lyle et al. 2009). Estimates in 2011/12 and 2012/13 were 31.9 and 15.2 t respectively for Striped Trumpeter and exceeded the commercial catch for the species in both years (Fig. 21.3A). These catch estimates do not fully represent catches by charter boats. Interestingly, in recent years good numbers of large Striped Trumpeter have been found in waters off Portland in western Victoria and have become increasingly targeted by recreational fishers.




Figure 21.1 Striped Trumpeter length composition (1999–2012) from commercial catches. n = sample size. Dotted line indicates the minimum legal size limit and * indicate years when the minimum size limit was changed. Length is fork length.




Figure 21.1 - continued

Table 21.1 Annual commercial catches of Striped Trumpeter (t) south of latitude 39 12'S. Data based on Tasmanian (General Fishing return), Victorian and Commonwealth catch returns.

Year	Catch (t)			Combined
	Tasmanian	Victoria	Commonwealth	
1990/91	74.5	37.1		111.6
1991/92	58.2	36.8		95.0
1992/93	52.7	19.8		72.5
1993/94	56.5	16.0		72.5
1994/95	72.4	14.6		87.0
1995/96	60.3			60.3
1996/97	79.7		0.7	80.4
1997/98	75.4		5.7	81.1
1998/99	98.4		8.9	107.4
1999/00	86.3		14.5	101.8
2000/01	41.2		7.5	49.6
2001/02	40.0		4.8	44.8
2002/03	36.8		3.2	40.0
2003/04	36.8		3.7	40.5
2004/05	24.0		2.2	26.2
2005/06	19.1		4.7	23.8
2006/07	18.8		3.5	22.3
2007/08	13.1		3.0	16.1
2008/09	10.5		2.8	13.3
2009/10	10.0		2.3	12.8
2010/11	15.0		4.8	19.8
2011/12	15.8		5.4	21.2
2012/13	12.2		0.8	13.0
2013/14	8.0		2.3	10.3




Figure 21.2 Striped Trumpeter age composition (1999–2012) from commercial catches. n = sample size (year 2009 to 2012 on following page).




Figure 21.2 (continued)

Apart from the north coast, Striped Trumpeter catches have historically been reported from all areas around the state (Figs. 21.3A and 21.4). Catches declined strongly between 1999/2000 and 2000/01. Since 2009/10, catches in the north-west coast have increased to match those of the south-east, with the majority of catches stemming from the west coast in 2013/14 (Figs. 21.3A and 21.4).

The observed catch trends reflect the influence of especially strong year classes that entered the fishery between 1995/96 and 1997/98. Increased graball catches in 1998/99 followed by a decline suggest that the 1996 year-class, which would have recruited to the inshore gillnet fishery in 1998/99, may also have been relatively strong. The subsequent decline in graball catches presumably reflects the movement of these relatively strong year-classes offshore but also suggests that there has been limited recruitment in recent years. Industry representatives also suggest that the trip limit of 250 kg introduced in 2000 provided a strong disincentive for some operators to target the species and may have contributed to the fall in drop-line and handline catches and effort since 2000/01.

Fishing effort increased during the latter part of the 1990s, presumably linked to the increased availability of Striped Trumpeter (Fig. 21.3B). Subsequently, effort for all methods has declined, with handline fishing remaining the dominant method. Fishing activity has been focussed mainly on the south-east and west coasts and to a lesser extent off the east and north-east (Fig. 21.3B).

Recent gear-specific trends of catch rates (including 2013/14) are based on small catches and may not be informative about availability (Fig. 21.3C). Graball catch rates increased steadily up until 2003/04, despite declining catches during the latter half of the period. The sharp fall in graball catch rates since 2004/05 may have been influenced by the minimum size limit increase that took effect during 2004.

Striped Trumpeter





Figure 21.3 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. HL= hand line, GN= graball net, DL = drop-line, SEC= south-east coast, EC= east coast, NEC= north-east coast, WC = west coast.

A) Catch



B) Effort




Figure 21.4 (A) Striped Trumpeter catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (80.4 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (52.7 t)	Yes	42.4 t (80.5%)
	• Catch increases by > 30% from previous year (>16.9 t)	No	
	• Catch decreases by > 30% from previous year (<9.1 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>1695.1 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0171 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (79.7 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (23.9 t)	Yes	13.6 t (57%)
	• Catch variation from the previous year above the greatest inter-annual increase from the reference period (21.1 t)	No	
	• Catch variation from the previous year above the greatest inter-annual decrease from the reference period (49.5 t)	No	
	• Latest recreational catch estimate > recreational catch estimate from the reference period (19.6 t)	Yes	2.3 t (7.8%)
	• Proportion of recreational catch to total catch > previous proportion estimate (61.1% in 2011/12%)	No	
Biomass	• CPUE < 3 rd lowest CPUE value from the reference period (0.0210 t/days fished)	No	
	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0034)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	No	
	• Significant numbers of unhealthy fish landed	No	

Stock status

TRANSITIONAL-RECOVERING

The sharp decline in commercial catches since 2000/01 gave rise to concern about the status of Striped Trumpeter stocks. As suggested in previous assessments, strong recruitment variability could result in marked variation in population size, especially if there is a prolonged period of poor recruitment. Recent age composition data provide some encouraging signs

regarding new recruits entering the fishery, although the relative strength of the new cohorts (5–6 year olds) is unclear. These fish are primarily immature and thus have yet to contribute to the spawning stock.

Research undertaken during 2010 confirmed that the current minimum size limit is well below the size at maturity (>60 cm TL cf minimum size limit of 50 cm TL) and that with the increasing proportion of small fish in the catch due to the recent recruitment events there is potential for growth overfishing if fishing pressure were to increase dramatically. However, as noted in previous assessments the 250 kg trip limit appears to have reduced the incentive for commercial fishers to target Striped Trumpeter. By contrast, growing interest from the recreational sector suggests that recreational catches represent an increasingly significant proportion of total fishing mortality. While the implementation of an on-water catch limit of four fish for recreational fishers may help to constrain recreational harvest, the presence of small legal sized fish may provide an incentive for some high grading as fishers seek to maximise the weight of their catch under the reduced bag limit.

Catches reported in Commonwealth returns in recent years have averaged about 3 t per year, though industry reports suggest that these figures may be significantly underestimated. There is an urgent need to ensure that catch and effort information are comprehensive and approaches have been made to the Commonwealth to this end. It should be noted, however, that AFMA Fisheries Compliance Officers do monitor the unloading of Commonwealth vessels at times and do carry out random vessel inspections.

A spawning season closure during September and October, when fish are particularly vulnerable to capture, was introduced in 2009 for Tasmanian commercial fishers. Fishing trials have confirmed that while there may be some variability in the timing of spawning around the state, the closure encompasses the bulk of the spawning season in all areas.


The influx of 4–6 year olds in the last 4 years provides positive signs of population recovery after a prolonged period of low recruitment, which has led to the stock status of transitional-recovering. The relative strength of these year classes will need to be monitored over time to determine how much they contribute to population rebuilding. Aligning the size limit with size at maturity, thereby allowing fish to spawn before they become vulnerable to capture, would almost certainly increase spawning biomass and enhance recruitment potential.

22. Wrasse

Blue-throated Wrasse *Notolabrus tetricus*

Purple Wrasse *Notolabrus fuciola*

STOCK STATUS	NOT ASSESSED
Both species are targeted as part as a live fishery. Recent ban on the preferred bait species in fish traps has led to a decline in participation in this fishery.	
IMPORTANCE	Key
STOCK(S)	Tasmanian Scalefish Fishery
INDICATOR(S)	Catch, effort and CPUE trends



Notolabrus tetricus
Source: DPIPWE (by Peter Gouldthorpe)

Species biology

Parameters	Estimates	Source
Habitat	<ul style="list-style-type: none"> <u>Bluethroat Wrasse</u>: Sheltered and exposed reefs, from 1 to 160m depth. <u>Purple Wrasse</u>: Predominantly on exposed reefs up to 90m depth. 	Edgar (2008)
Distribution	<ul style="list-style-type: none"> <u>Bluethroat Wrasse</u>: from Sydney (New South Wales) to Ceduna (South Australia). <u>Purple Wrasse</u>: New Zealand and Australian waters, from southern New South Wales to Kangaroo Island (South Australia). 	Edgar (2008)
Diet	<ul style="list-style-type: none"> Both species consume a range of molluscs and crustaceans. 	Shepherd & Clarkson (2001) Denny & Schiel (2001)
Movement and stock structure	<ul style="list-style-type: none"> <u>Bluethroat Wrasse</u>: site-attached. Females have overlapping home ranges and males are territorial, at least during the reproductive season. No emigration of individuals between reefs has been noted for this species. <u>Purple Wrasse</u>: site-attached, with no evidence of territorial behaviour. Movements between reefs is limited and likely to be restricted to a small proportion of the population. No information on the stock structure for any species 	Barrett (1995b)
Natural mortality	<ul style="list-style-type: none"> Low adult mortality for both species. Estimated at $M = 0.2$ for blue-throated Wrasse. 	Smith et al. (2003) Barrett (1995a)
Maximum age	<ul style="list-style-type: none"> <u>Bluethroat Wrasse</u>: 11 years <u>Purple Wrasse</u>: up to 24 years 	Barrett (1995a) Welsford (2003)

Growth	<ul style="list-style-type: none"> • Bluethroat Wrasse: From 6 months onwards, growth (males and females confounded) follows a Von Bertalanffy growth function with $L_{\infty} = 36.12$, $k = 0.2$ and $t_0 = -0.35$. Length-weight relationship was set at $W = 0.0545L^{2.7157}$ (both sexes) where W is weight (g) and L is the fork length (cm). • Purple Wrasse: From 2 years onwards, growth (males and females confounded) follows a Von Bertalanffy growth function with $L_{\infty} = 44.7$, $k = 0.085$ and $t_0 = -3.23$. Length-weight relationship (both sexes) was set at $W = 0.0161L^{3.0407}$ where W is weight (g) and L is the fork length (cm). 	Welsford (2003) Barrett (1995a) Unpublished data
Maturity	<ul style="list-style-type: none"> • Bluethroat Wrasse: protogynous hermaphrodite (i.e. developing into female first before changing to male) with sex change happening between 27 and 32 cm in Tasmania. However, not all individuals undergo a sex inversion. Size at 50% maturity reached at 29.89 cm for females (corresponding to around 8 years old). Batch fecundity unknown. • Purple Wrasse: gonochoristic species (i.e. sex is fixed at maturity). Size at 50% maturity reached at 18.41 cm for females (corresponding to around 3 years old). Batch fecundity is estimated at 74500 ± 34900 eggs/kg. 	Barrett (1995a) Hardwood & Lokman (2006) Unpublished data
Spawning	<ul style="list-style-type: none"> • The spawning season for both species extends from August to January. 	Barrett (1995a)
Early life history	<ul style="list-style-type: none"> • Bluethroat Wrasse: planktonic larval duration ranges from 44 to 66 days. • Purple Wrasse: planktonic larval duration ranges from 40 to 87 days. • Settlement on the reefs at around 10.4mm for both species. 	Welsford (2003)
Gillnet post release survival	<ul style="list-style-type: none"> • Moderate – High: 83 – 59% depending on gillnet soak duration 	Lyle et al (2014a)

Background

Of the several species of Wrasse occurring in Tasmanian waters, Purple Wrasse (*Notolabrus fucicola*) and Blue-throat Wrasse (*N. tetricus*) are the main species taken commercially. Wrasse are targeted for live fish markets as well as being sold as dead product and utilised as bait for rock lobster (bait usage is likely to be under-reported). Fish marketed live are distinguished in the logbooks, and live Wrasse have accounted for over 90% of the total reported catch since 2001/02. Thus, trends in the live-fish fishery will ultimately be reflected in overall production levels. The two species of Wrasse have only been distinguished in catch returns since 2007. While there is an apparent market preference for Blue-throat Wrasse, Purple Wrasse are more robust for live handling.

FISHING METHODS	Fish trap and hand line
MANAGEMENT METHODS	Input control: <ul style="list-style-type: none"> • Gear licence (Scalefish fishing licence) • Species licence (Wrasse licence) • Rock lobster licence (for bait only)

MAIN MARKET	Output control: <ul style="list-style-type: none"> • Legal size: 30 cm. • Possession limit of 10 (recreational fishers) • Interstate (live trade) and local (bait and food).
--------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Current assessment

Catch, effort and CPUE

Wrasses are targeted all around Tasmania, the west coast being the least fished area (Fig. 22.2). Wrasse catches fluctuated between 75–110 t between 1995/96 and 2007/08, peaking at 113 t in 2006/07 (Fig. 22.1A). Reported catches fell in 2008/09 to just 68.2 t (42 t of Blue-throat Wrasse, 26 t of Purple Wrasse, 0.2 t undefined Wrasse) and have remained at a similar level since, reaching 62.4 t (49.9 t Blue-throat, 12.5 t Purple, 0.4 t undefined Wrasse) in 2013/14. The decline in catches since 2006/07 is largely due to a decline in the use of fish traps as a result of the prohibition on the use of abalone gut as bait. This prohibition was a response to the appearance of the abalone viral ganglioneuritis (AGV) in Victoria and has forced fishers to seek alternative but less effective baits. Catches, effort and catch rates subsequently declined for fish traps, particularly in the south-east (Fig. 22.1) where fish traps were the dominant fishing method. By contrast, hand line catch and effort has remained relatively stable since that time.

With Blue-throat Wrasse being more susceptible to line methods and Purple Wrasse more vulnerable to trap capture, Blue-throat Wrasse are now taken in larger quantities in the live fishery. Gillnets account for the bulk of the remaining catch (< 5 t) but because survival in nets is poor, graball caught Wrasse are rarely marketed live.

Recreational catches were estimated at 13.6 t in 2000/01 (Lyle 2005) and 10.3 t in 2007/08 (Lyle et al. 2009) and 6.4 t in 2012/13 (Lyle et al. 2014b), which represent around 10% of the total Tasmanian catches for these species in each instance. Further, Blue throated wrasse are a reasonably common by-catch of recreational gillnet fishers and a recent study showed that they have a moderate to low post-release survival, particularly when gillnets are deployed for > 4 hours (Lyle et al. 2014a).

Catch rate trends imply that Wrasse stocks have not been impacted significantly by the fishery until recent years. Catch rates for both gear types stabilised at moderate levels after 2008/09, with slight increases in the last two years. Catch rates for fish traps peaked in 2004/05, but have since declined, largely influenced by changes in permissible bait types (Fig. 22.1C). There is, nonetheless evidence that exploitation rates for legal-sized Purple Wrasse on some east-coast reefs are extremely high (Ewing 2004). The broad-scale analyses presented here are, however, generally insensitive to changes in abundance at the level of individual reefs at which the fishery impacts the populations. Marked regional shifts of effort have occurred in the fishery over the years and may have also masked localised depletions, with fishers moving to new or lightly fished areas to maintain catches.




Figure 22.1 A) Annual commercial catch (t) by gear (left) and region (right), and best estimates of recreational catches (blue squares). B) Commercial effort by method based on gear units (left) and day fished (right) relative to 1995/96. C) Commercial catch per unit effort (CPUE) based on weight per gear unit (left) and weight per day fished (right) relative to 1995/96. HL= hand line, FP= fish trap, SEC= south east coast, EC= east coast, NEC= north-east coast, NWC = north-west coast.




Figure 22.2 (A) Wrasse catches (t) and (B) effort (days) by fishing blocks averaged from 2008/09 to 2012/13 (left) and during 2013/14 (right).

Reference points

Performance indicators	Current reference points	Breached?	By how much?
Catch	• Catch > higher catch from the 1990/91 to 1997/98 range (178 t)	No	
	• Catch < lowest catch from the 1990/91 to 1997/98 range (57.2 t)	No	
	• Catch increases by > 30% from previous year (>93.1 t)	No	
	• Catch decreases by > 30% from previous year (<50.1 t)	No	
Effort trend	• Effort >10% of highest level from 1995/96 to 1997/98 (>4464 days fished)	No	
Catch rates trends	• Catch rate < 80% of lowest levels from 1995/96 to 1997/98 (<0.0093 t/days fished)	No	

Performance indicators	Proposed reference points	Breached?	By how much?
Fishing mortality	• Catch > 3 rd highest catch value from the reference period (100.1 t)	No	
	• Catch < 3 rd lowest catch value from the reference period (83.4 t)	Yes	20.8 t (24.9%)
	• Catch variation from the previous year above the greatest inter-annual increase from the reference period (26.7 t)	No	
	• Catch variation from the previous year above the greatest inter-annual decrease from the reference period (25.3 t)	No	
	• Latest recreational catch estimate > recreational catch estimate from the reference period (13.6 t)	No	
	• Proportion of recreational catch to total catch > previous proportion estimate (13.1% in 2007/08)	No	
Biomass	• CPUE < 3 rd lowest CPUE value from the reference period (0.0135 t/days fished)	No	
	• Rate of CPUE decline over last 3 years is greater than the largest 3-year CPUE decline during the reference period (-0.0014)	No	
Stock stress	• Significant change in the size/age composition of commercial catches	Not assessed	
	• Significant numbers of unhealthy fish landed	No	

Stock status

NOT ASSESSED

The reference for the lowest catch was breached, which most likely reflects the changes in fishing behaviour (away from traps) rather than a stock issue since catch rates remain relatively high.

While input controls (limited entry) have capped participation in the live Wrasse fishery, there is still a substantial level of latent effort. Increasing catches up to 2006/07 reflected a strong interest in the species, and there were concerns that these levels were not sustainable. Under present arrangements, there is potential for localised depletions of legal-sized Wrasse, especially if effort becomes concentrated in particular regions.

The minimum size limit provides good protection (several years after reaching the size at maturity) for the spawning stock of Purple Wrasse and for female Blue-throat Wrasse. The limit does not, however, provide the same level of protection for male Blue-throat Wrasse because males are derived through sex change from mature females, typically at sizes after they have entered the fishery. This, along with the fact that males are strongly site attached and have higher catchability (being more aggressive than females), suggests that they are vulnerable to fishing. In extreme situations it is possible that localised heavy fishing pressure could result in 'sperm shortage' that could affect spawning success even though there may be a robust population of mature (sub-legal size) females present. The removal of the maximum size limit for Wrasse in 2004 may have exacerbated this potential problem. However there are no clear indications of spawning stock shortages in Tasmania.

The banning of the preferred bait for trap fishing in 2008 has dramatically changed fisher's behaviour and fish traps are now rarely used. This has resulted in Blue-throat Wrasse contributing a greater proportion of the catch than beforehand. Industry reported that there is no acceptable substitute for abalone guts (i.e. no other bait yields the same return) and that trap fishing has become less viable. This is supported by catch rate data in Figure 22.1. As a consequence trap fishing effort has markedly decreased along with line fishing, which was previously an associated activity.

Due to the uncertainties highlighted above, there is insufficient information available to confidently classify the status of the two species.

References

- AFMA. 2008. Blue warehou (*Seriolella brama*) stock rebuilding strategy: Southern and Eastern Scalefish and Shark Fishery (SESSF) - December 2008. Australian Fisheries Management Authority, Canberra.
- AFMA. 2011. Blue warehou (*Seriolella brama*) stock rebuilding strategy: Southern and Eastern Scalefish and Shark Fishery (SESSF) - Revised July 2011. Australian Fisheries Management Authority, Canberra.
- AFMA. 2012. Blue warehou (*Seriolella brama*) stock rebuilding strategy: Southern and Eastern Scalefish and Shark Fishery (SESSF) - Revised April 2012. Australian Fisheries Management Authority, Canberra.
- AFMA. 2014. Blue warehou (*Seriolella brama*) Stock Rebuilding Strategy. Australian Fisheries Management Authority, Canberra, ACT.
- Annala, J. H. 1994. Report from the Fishery Assessment Plenary, May 1994: stock assessments and yield estimates (unpublished). MAF Fisheries Greta Point library, Wellington
- Ayling, G. M., K. C. Wilson, and D. A. Ratkowsky. 1975. Sand flathead (*Platycephalus bassensis*), an indicator species for mercury pollution in Tasmanian waters. *Marine Pollution Bulletin* **6**:142-144.
- Bani, A. 2005. Temporal and spatial variability of the life history characteristics of sand flathead, *Platycephalus bassensis*. University of Tasmania, Hobart.
- Bani, A. and N. A. Moltschanivsky. 2008. Spatio-temporal variability in reproductive ecology of sand flathead, *Platycephalus bassensis*, in three Tasmanian inshore habitats: potential implications for management. *Journal of Applied Ichthyology* **24**:555-561.
- Barnes, L. M., C. A. Gray, and J. E. Williamson. 2011. Divergence of the growth characteristics and longevity of coexisting Platycephalidae (Pisces). *Marine and Freshwater Research* **62**:1308-1317.
- Barrett, N. S. 1995a. Aspects of the biology and ecology of six temperate reef fishes (Families: Labridae and Monacanthidae). University of Tasmania, Hobart.
- Barrett, N. S. 1995b. Short- and long-term movement patterns of six temperate reef fishes (Families Labridae and Monacanthidae). *Marine and Freshwater Research* **46**:853-860.
- Bertoni, M. 1995. The reproductive biology and feeding habits of the snook, *Sphyrna novaehollandiae*, in South Australian waters. *Southern Fisheries* **3**:34-35.
- Blackburn, M. 1960. A study of condition (weight for length) of Australian barracouta, *Thyrsites atun* (Euphrasen). *Australian Journal of Marine and Freshwater Research* **11**:14-41.
- Blackburn, M. and P. E. Gartner. 1954. Populations of barracouta, *Thyrsites atun* (Euphrasen), in Australian waters. *Australian Journal of Marine and Freshwater Research* **5**:411-468.
- Bruce, B. D., K. Evans, C. A. Sutton, J. W. Young, and D. M. Furlani. 2001a. Influence of mesoscale oceanographic processes on larval distribution and stock structure in jackass morwong (*Nemadactylus macropterus*: Cheilodactylidae). *ICES Journal of Marine Science* **58**:1072-1080.
- Bruce, B. D., F. J. Neira, and R. W. Bradford. 2001b. Larval distribution and abundance of blue and spotted warehou (*Seriolella brama* and *S. punctata*: Centrolophidae) in south-eastern Australia. *Marine and Freshwater Research* **52**:631-639.
- Bulman, C., F. Althaus, X. He, N. J. Bax, and A. Williams. 2001. Diets and trophic guilds of demersal fishes of the south eastern Australian shelf. *Marine and Freshwater Research* **52**:537-548.
- Bulman, C., S. Condie, J. Findlay, B. Ward, and J. Young. 2008. Management zones from small pelagic fish species stock structure in southern Australian waters. FRDC report 2006/076. CSIRO, Hobart.
- Burchmore, J. J., D. A. Pollard, M. J. Middleton, J. D. Bell, and B. C. Pease. 1988. Biology of four species of whiting (Pisces: *Sillaginidae*) in Botany Bay, New South Wales. *Australian Journal of Marine and Freshwater Research* **39**:709-727. .

- Butcher, A. R. and A. R. Hagedoorn. 2003. Age, growth and mortality estimates of stout whiting, *Sillago robusta* Stead (Sillaginidae), from Southern Queensland, Australia. *Asian Fisheries Science* **16**: 215-228.
- Buxton, C. D., J. M. Semmens, E. Forbes, J. M. Lyle, N. S. Barrett, and M. J. Phelan. 2010. Spatial management of reef fisheries and ecosystems: understanding the importance of movement. Tasmanian Aquaculture and Fisheries Institute, Hobart.
- BWAG. 1998. Stock assessment report: Blue warehou 1998. Australian Fisheries Management Authority, Canberra.
- Chubb, C. F., I. C. Potter, C. J. Grant, R. C. J. Lenanton, and J. Wallace. 1981. Age structure, growth rates and movements of sea mullet, *Mugil cephalus* L. and yellow-eye mullet, *Aldrichetta forsteri* (Valenciennes), in the Swan-Avon River System, Western Australia. *Australian Journal of Marine and Freshwater Research* **32**:605-628.
- Coleman, N. and M. Mobley. 1984. Diets of commercially exploited fish from Bass Strait and adjacent Victorian waters, South-eastern Australia. *Australian Journal of Marine and Freshwater Research* **35**:549-560.
- Crawford, C. M. 1984. An ecological study of Tasmanian flounder. University of Tasmania, Hobart.
- Crawford, C. M. 1986. Development of eggs and larvae of the flounders *Rhombosolea tapirina* and *Ammotretis rostratus* (Pisces: Pleuronectidae). *Journal of Fish Biology* **29**:325-334.
- Curtis, T. D. and J. S. Shima. 2005. Geographic and sex specific variation in growth of yellow eyed mullet, *Aldrichetta forsteri*, from estuaries around New Zealand. *New Zealand Journal of Marine and Freshwater Research* **39**:1277-1285.
- Denny, C. M. and D. R. Schiel. 2001. Feeding ecology of the banded wrasse *Notolabrus fucicola* (Labridae) in southern New Zealand: prey items, seasonal differences, and ontogenetic variation. *New Zealand Journal of Marine and Freshwater Research* **35**:925-933.
- Dixon, P. I. 1987. Stock identification and discrimination of commercially important whittings in Australian waters using genetic (FIRTA 83/16): Final Report. University of New South Wales, Centre for Marine Science.
- DPIF. 1998. Scalefish Fishery: Policy Document. Tasmanian Department of Primary Industry and Fisheries, Hobart.
- Dunning, M. C. 1998. Zoogeography of arrow squids (Cephalopoda: Ommastrephidae) in the Coral and Tasman Seas, southwest Pacific. Pages 435-453 in N. A. Voss, M. Vecchione, R. B. Toll, and M. J. Sweeney, editors. *Systematics and Biogeography of Cephalopods*. Smithsonian Institution Press, Washington D.C.
- Dunning, M. C. and E. C. Förch. 1998. A review of the systematics, distribution and biology of arrow squids of the genus *Nototodarus* Pfeffer, 1912 (Cephalopoda, Ommastrephidae). Pages 393-404 in N. A. Voss, M. Vecchione, R. B. Toll, and M. J. Sweeney, editors. *Systematics and Biogeography of Cephalopods*. Smithsonian Institution Press, Washington D.C.
- Edgar, G. D. 2008. Australian marine life: the plants and animals of temperate waters. New Holland Publishers, Sydney.
- Edgar, G. J. 1997. Australian marine life: the plants and animals of temperate waters. Reed Books, Melbourne.
- Edgar, G. J., N. S. Barrett, and A. J. Morton. 2004. Patterns of fish movement on eastern Tasmanian rocky reefs. *Environmental Biology of Fishes* **70**:273-284.
- Elliott, N. G. and R. D. Ward. 1994. Enzyme variation in jackass morwong, *Nemadactylus macropterus* (Schneider 1801) (Teleostei: Cheilodactylidae) from Australian and New Zealand waters. *Australian Journal of Marine and Freshwater Research* **45**:51-67.
- Erwing, G., P., J. M. Lyle, R. Murphy, J. M. Kalish, and P. E. Ziegler. 2007. Validation of age and growth in a long-lived temperate reef fish using otolith structure, oxytetracycline and bomb radiocarbon methods. *Marine and Freshwater Research* **58**:944-955.
- Ewing, G., P. 2004. Spatial and temporal variation in growth and age composition of the temperate wrasse *Notolabrus fucicola* in Tasmanian waters. MSc thesis. University of Tasmania, Hobart.

- Fairbridge, W. S. 1951. The New South Wales tiger flathead, *Neoplatycephalus macrodon* (Ogilby). I. Biology and age determination. Australian Journal of Marine and Freshwater Research **2**:117-178.
- Ferguson, G. 2006. Fisheries biology of the greenback flounder *Rhombosolea tapirina* (Günther 1862) (Teleostei: Pleuronectidae) in South Australia. SARDI Aquatic Sciences Publication No. RD06/0008-1. South Australian Research and Development Institute (Aquatic Sciences), Adelaide.
- FishBase. 2013. Monacanthidae. Accessed 30/09/2013.
- Flood, M., I. Stobutzki, J. Andrews, C. Ashby, G. Begg, R. Fletcher, C. Gardner, L. Georgeson, S. Hansen, K. Hartmann, P. Hone, P. Horvat, L. Maloney, B. McDonald, A. Moore, A. Roelofs, K. Sainsbury, T. Saunders, T. Smith, C. Stewardson, J. Stewart, and B. Wise. 2014. Status of key Australian fish stocks reports 2014. Fisheries Research and Development Corporation, Canberra, ACT.
- Flood, M., I. Stobutzki, J. Andrews, G. Begg, R. Fletcher, C. Gardner, J. Kemp, A. Moore, A. O'Brien, R. Quinn, J. Roach, K. R. Rowling, K. Sainsbury, T. Saunders, T. Ward, and M. Winning. 2012. Status of key Australian fish stocks reports 2012. Fisheries Research and Development Corporation, Canberra.
- Francis, M. 2001. Coastal fishes of New Zealand. Reed Publishing, Auckland.
- Frimodt, C. 1995. Multilingual illustrated guide to the world's commercial coldwater fish. Wiley-Blackwell, Oxford.
- Gavrilov, G. M. and N. P. Markina. 1979. The feeding ecology of fishes of the genus *Seriolella* (fam. Nomeidae) on the New Zealand plateau. Journal of Ichthyology **19**:128-135.
- Georgeson, L., I. Stobutzki, and R. Curtotti. 2014. Fishery status reports 2013–14. Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra, ACT.
- Godfriaux, B. L. 1974. Food of tarakihi in western Bay of Plenty and Tasman Bay, New Zealand. New Zealand Journal of Marine and Freshwater Research **8**:111-153.
- Goldworthy, S. D., C. Bulman, X. He, J. Larcombe, and C. Littnan. 2003. Trophic interactions between marine mammals and Australian fisheries: ecosystem approach. Pages 62-99 in N. Gales, M. Hindell, and R. Kirkwood, editors. Marine mammals: fisheries, tourism and management issues. CSIRO Publishing, Collingwood, Victoria.
- Gomon, M., D. Bray, and R. Kuitert. 2008. Fishes of Australia's southern coast. Reed New Holland.
- Gorman, T. B. S. 1962. Yellow-eyed mullet *Aldrichetta forsteri* Cuvier and Valenciennes in Lake Ellesmere.
- Grant, C. J., T. R. Cowper, and D. D. Reid. 1978. Age and growth of snoek, *Leionura atun* (Euphrasen), in south-eastern Australian waters. Australian Journal of Marine and Freshwater Research **29**:435-444.
- Grant, E. M. 1991. Grant's Fishes of Australia. E.M. Grant Pty. Ltd., Queensland.
- Grewe, P. M., A. J. Smolenski, and R. D. Ward. 1994. Mitochondrial DNA variation in jackass morwong, *Nemadactylus macropterus* (Teleostei: Cheilodactylidae) from Australian and New Zealand waters. Canadian Journal of Fisheries and Aquatic Sciences **51**:1101-1109.
- Hardwood, N. J. and M. P. Lokman. 2006. Fecundity of banded wrasse (*Notolabrus fucicola*) from Otago, Southern New Zealand. New Zealand Journal of Marine and Freshwater Research **40**:467-476.
- Harries, D. N. and R. L. Croome. 1989. A review of past and present inshore gill netting in Tasmania with particular reference to the bastard trumpeter, *Latridopsis forsteri* Castelnau. papers and Proceedings of the Royal Society of Tasmania **123**:97-110.
- Harries, D. N. and P. S. Lake. 1985. Aspects of the biology of inshore populations of Bastard Trumpeter, *Latridopsis forsteri* (Castleneau, 1872) in Tasmanian waters. Tasmanian Fisheries Research **27**:19-43.
- Hartmann, K. and J. M. Lyle. 2011. Tasmanian Scalefish Fishery-2009/10. Tasmanian Aquaculture and Fisheries Institute, Hobart.

- Henry, G. W. and J. M. Lyle. 2003. National recreational and indigenous fishing survey. Final report to FRDC. Project No 99/158. Tasmanian Aquaculture and Fisheries Institute, Hobart.
- Hobday, D. K. and J. W. J. Wankowski. 1987. School whiting *Sillago bassensis flindersi*: reproduction and fecundity in eastern Bass Strait, Australia. Internal Report 153. Victorian Department of Conservation, Forests and Lands. Fisheries Division.
- Hurst, R. J., S. L. Ballara, and D. MacGibbon. 2012. Fishery characterisation and standardised CPUE analyses for barracouta, *Thyrsites atun*, (Euphrasen, 1791) (Gempylidae), 1989–90 to 2007–08. New Zealand Fisheries Assessment Report 2012/12 NIWA, Wellington.
- Hutchinson, W. 1993. The reproductive biology and induced spawning of striped trumpeter, *Latris lineata*. University of Tasmania, Hobart.
- Jackson, G. D., B. McGrath Steer, S. Wotherspoon, and A. J. Hobday. 2003. Variation in age, growth and maturity in the Australian arrow squid *Nototodarus gouldi* over time and space-what is the pattern? Marine Ecology Progress Series **264**:57-71.
- Jackson, G. D., R. K. O'Dor, and Y. Andrade. 2005a. First tests of hybrid acoustic/archival tags on squid and cuttlefish. Marine and Freshwater Research **56**:425-430.
- Jackson, G. D., S. Wotherspoon, and B. L. McGrath-Steer. 2005b. Temporal population dynamics in arrow squid *Nototodarus gouldi* in southern Australian waters. Marine Biology **146**:975-983.
- Jenkins, G. P. 1986. Composition, seasonality and distribution of ichthyoplankton in Port Phillip Bay, Victoria. Australian Journal of Marine and Freshwater Research **37**:507-520.
- Jones, G. K. 1990. Growth and mortality in a lightly fished population of Garfish (*Hyporhamphus melanochir*), in Baird Bay, South Australia. Transactions of the Royal Society of South Australia **114**:37-45.
- Jones, G. K., Q. Ye, S. Ayvazian, and P. Coutin. 2002. Fisheries ecology and habitat ecology of southern sea Garfish (*Hyporhamphus melanochir*) in southern Australian waters. South Australian Research and Development Institute.
- Jordan, A. R. 2001a. Age, growth and spatial and interannual trends in age composition of jackass morwong, *Nemadactylus macropterus*, in Tasmania. Marine and Freshwater Research **52**:641-660.
- Jordan, A. R. 2001b. Reproductive biology, early life-history and settlement distribution of sand flathead (*Platycephalus bassensis*) in Tasmania. Marine and Freshwater Research **52**:589-601.
- Jordan, A. R., D. M. Mills, G. Ewing, and J. M. Lyle. 1998. Assessment of inshore habitats around Tasmania for life-history stages of commercial finfish species. Tasmanian Aquaculture and Fisheries Institute, Hobart.
- Kailola, P. J., M. J. Williams, P. C. Stewart, R. E. Reichelt, A. McNee, and C. Grieve. 1993. Australian fisheries resources. Bureau of Resource Sciences, Canberra.
- Klaer, N. 2010. Tiger flathead (*Neoplatycephalus richardsoni*) stock assessment based on data up to 2009. CSIRO, Hobart.
- Klumpp, D. W. and P. D. Nichols. 1983. Nutrition of the southern sea Garfish (*Hyporhamphus melanochir*): gut passage rate and daily consumption of two food types and assimilation of seagrass components. Marine Ecology Progress Series **12**:207-216.
- Knuckey, I. A. and K. P. Sivakumaran. 2001. Reproductive characteristics and per-recruit analyses of blue warehou (*Seriolella brama*): implications for the south-east fishery of Australia. Marine and Freshwater Research **52**:575-587.
- Last, P. R., E. O. G. Scott, and F. H. Talbot. 1983. Fishes of Tasmania. Tasmanian Fisheries Development Authority Hobart.
- Lyle, J. M. 1999. Licensed recreational fishing and an evaluation of recall biases in the estimation of recreational catch and effort. Final Report to the Marine Recreational Fishing Council Tasmanian Aquaculture and Fisheries Institute, Hobart.
- Lyle, J. M. 2000. Assessment of the licensed recreational fishery of Tasmania (Phase 2). Final Report to FRDC. Project 96/161. Tasmanian Aquaculture and Fisheries Institute Hobart.
- Lyle, J. M. 2005. 2000/01 survey of recreational fishing in Tasmania. Tasmanian Aquaculture and Fisheries Institute, Hobart.

- Lyle, J. M., J. D. Bell, B. M. Chuwen, N. S. Barrett, S. R. Tracey, and C. D. Buxton. 2014a. Assessing the impacts of gillnetting in Tasmania: Implications for by-catch and biodiversity. Institute for Marine and Antarctic Studies, Hobart, Tasmania.
- Lyle, J. M. and W. B. Ford. 1993. Review of trawl research 1979–1987, with summaries of biological information for the major species, with summaries of biological information for the major species. Technical Report 46. Department of Sea Fisheries, Tasmania, Hobart.
- Lyle, J. M. and A. R. Jordan. 1999. Tasmanian scalefish fishery assessment- 1998. Tasmanian Aquaculture and Fisheries Institute, Hobart.
- Lyle, J. M., K. Krusic-Golub, and A. K. Morison. 2000. Age and growth of jack mackerel and the age structure of the jack mackerel purse seine catch. FRDC project no 1995/034. TAFI, Hobart.
- Lyle, J. M., N. Moltschaniwskyj, A. J. Morton, I. W. Brown, and D. Mayer. 2007. Effects of hooking damage and hook type on post-release survival of sand flathead (*Platycephalus bassensis*). *Marine and Freshwater Research* **58**:445-453.
- Lyle, J. M. and R. Murphy. 2001. Long distance migration of striped trumpeter. *Fishing Today* **14**:6.
- Lyle, J. M., K. E. Stark, and S. R. Tracey. 2014b. 2012-13 survey of recreational fishing in Tasmania. Institute for Marine and Antarctic Studies, Hobart, Tasmania.
- Lyle, J. M. and S. R. Tracey. 2012. Recreational gillnetting in Tasmania – an evaluation of fishing practices and catch and effort. Report to FishWise. IMAS, Hobart.
- Lyle, J. M., S. R. Tracey, K. E. Stark, and S. Wotherspoon. 2009. 2007-08 survey of recreational fishing in Tasmania. Tasmanian Aquaculture and Fisheries Institute, Hobart.
- Machida, S. 1983. A brief review of the squid survey by Hoyo Maru No. 67 in southeast Australian waters in 1979/80. *Memoirs of the National Museum Victoria* **44**:291-295.
- May, J. L. and J. G. H. Maxwell. 1986. Trawl fish from temperate waters of Australia. CSIRO Division of Fisheries Research, Hobart.
- McCormick, M. I. 1989a. Reproductive ecology of the temperate reef fish *Cheilodactylus spectabilis* (*Pisces: Cheilodactylidae*). *Marine Ecology Progress Series* **55**:113-120.
- McCormick, M. I. 1989b. Spatio-temporal patterns in the abundance and population structure of a large temperate reef fish. *Marine Ecology Progress Series* **53**:215-225.
- McCormick, M. I. 1998. Ontogeny of diet shifts by a microcarnivorous fish, *Cheilodactylus spectabilis* : relationship between feeding mechanics, microhabitat selection and growth. *Marine Biology* **132**:9-20.
- Moltschaniwskyj, N. A. and G. T. Pecl. 2003. Small-scale spatial and temporal patterns of egg production by the temperate loliginid squid *Sepioteuthis australis*. *Marine Biology* **142**:509-516.
- Morison, A. K., I. A. Knuckey, C. A. Simpfendorfer, and R. C. Buckworth. 2012. 2011 Stock assessment summaries for the South East Scalefish and Shark Fishery. Southern and Eastern Scalefish and Shark Fishery Assessment Group, AFMA.
- Murphy, R. and J. M. Lyle. 1999. Impact of gillnet fishing on inshore temperate reef fished, with particular reference to banded morwong. Final report to FRDC, project No. 95/145. Tasmanian Aquaculture and Fisheries Institute, Hobart.
- Nakamura, I. and N. V. Parin. 1993. FAO Species Catalogue. Vol. 15. Snake mackerels and cutlassfishes of the world (families Gempylidae and Trichiuridae). An annotated and illustrated catalogue of the snake mackerels, snoeks, escolars, gemfishes, sackfishes, domine, oilfish, cutlassfishes, scabbardfishes, hairtails, and frostfishes known to date. FAO.
- Neira, F. J., A. G. Miskiewicz, and T. Trnski. 1998. Larvae of temperate Australian fishes: laboratory guide for larval fish identification. University of Western Australia Press.
- Nichols, D. S., D. Williams, G. A. Dunstan, P. D. Nichols, and J. K. Volkman. 1994. Fatty acid composition of Antarctic and temperate fish of commercial interest. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **107**:357-363.
- Norman, M. 2000. Cephalopods, a world guide. ConchBooks, Hackenheim.
- Norman, M. and A. Reid. 2000. A guide to squid, cuttlefish and octopuses of Australasia. CSIRO Publishing/The Gould League of Australia, Collingwood/Moorabbin

- O'Shea, S., K. S. Bolstad, and P. A. Ritchie. 2004. First records of egg masses of *Nototodarus gouldi* McCoy, 1888 (Mollusca: Cephalopoda: Ommastrephidae), with comments on eggmass susceptibility to damage by fisheries trawl. *New Zealand Journal of Zoology* **31**:161-166.
- O'Sullivan, D. and J. M. Cullen. 1983. Food of the Squid *Nototodarus gouldi* in Bass Strait. *Australian Journal of Marine and Freshwater Research* **34**:261-285.
- Parker, S. and D. Fu. 2011. Age composition of the commercial tarakihi (*Nemadactylus macropterus*) catch in quota management area TAR 2 in fishery year 2009-2010. *New Zealand Fisheries Assessment Report* 2011/59.
- Paul, L. J. 2000. *New Zealand fishes. Identification, natural history and fisheries.* Reed Publishing, Auckland.
- Paul, L. T. and P. R. Taylor. 1998. A summary of the biology, recreational and commercial landings, and stock assessment of yellow-eyed mullet, *Aldrichetta forsteri* (Cuvier and Valenciennes, 1836) (Mugilidae: Mugilidae). *New Zealand Fisheries Assessment Research Document* 98/17, NIWA, Wellington.
- Pecl, G. T. 2000. Comparative life-history of tropical and temperate *Sepioteuthis* squids in Australian waters. James Cook University, Townsville.
- Pecl, G. T. 2004. The *in situ* relationships between season of hatching, growth and condition in the southern calamary, *Sepioteuthis australis*. *Marine and Freshwater Research* **55**:429-438.
- Pecl, G. T. and N. A. Moltschanivsky. 2006. Life history of a short-lived squid (*Sepioteuthis australis*): resource allocation as a function of size, growth, maturation, and hatching season. *ICES Journal of Marine Science* **63**:995-1004.
- Pecl, G. T., N. A. Moltschanivskyj, S. R. Tracey, and A. R. Jordan. 2004. Inter-annual plasticity of squid life-history and population structure: ecological and management implications. *Oecologia (Berlin)* **139**:515-524.
- Pecl, G. T., S. R. Tracey, J. M. Semmens, and G. D. Jackson. 2006. Use of acoustic telemetry for spatial management of *Sepioteuthis australis*, a highly mobile inshore squid species. *Marine Ecology Progress Series* **328**:1-15.
- Richardson, B. J. 1982. Geographical distribution of electrophoretically detected protein variation in Australian commercial fishes. II: Jackass morwong, *Cheilodactylus macropterus* (Bloch and Schneider). *Australian Journal of Marine and Freshwater Research*, **33**: 927–931. **33**:927-931.
- Robinson, N., A. Skinner, L. Sethuraman, H. McPartlan, N. Murray, I. Knuckey, D. C. Smith, J. Hindell, and S. Talman. 2008. Genetic stock structure of blue-eye trevalla (*Hyperoglyphe antarctica*) and warehou (*Seriola lalandi* and *Seriola punctata*) in south-eastern Australian waters. *Marine and Freshwater Research* **59**:502-514.
- Rowling, K. R. 1994. Tiger flathead, *Neoplatycephalus richardsoni*. Pages 124-136 in R. D. J. Tilzey, editor. *The South East Fishery: a scientific review with reference to quota management.* Bureau of Resource Sciences, Australian Government Print Service, Canberra.
- Ruwald, F. P. 1992. Larval feeding trials with striped trumpeter, *Latris lineata*. in D. A. Hancock, editor. *Larval biology.* Australian Society for Fish Biology Workshop, Hobart, 20 August 1991. Bureau of Rural Resources Proceedings, AGPS, Canberra.
- Ruwald, F. P., L. D. Searle, and L. A. Oates. 1991. A preliminary investigation into the spawning and larval rearing of striped trumpeter, *Latris lineata*.
- Sahlqvist, P. and M. Skirtun. 2011. Southern Squid Fishery. in J. Woodhams, I. Stobutzki, S. Vieira, R. Curtotti, and G. Begg, editors. *Fisheries Status Reports 2010: status of fish stocks and fisheries managed by the Australian Government.* Australian Bureau of Agriculture and Resource Economics and Sciences, Canberra.
- Schnute, J. T. and L. J. Richards. 1990. A unified approach to the analysis of fish growth, maturity, and survivorship data. *Canadian Journal of Fisheries and Aquatic Sciences* **47**:24-40.
- Scott, T. D., C. J. M. Glover, and R. V. Southcott. 1974. *The marine and freshwater fishes of South Australia*, 2nd ed. Government Printer, South Australia.

- Shepherd, S. A. and P. S. Clarkson. 2001. Diet, feeding behaviour, activity and predation of the temperate blue-throated wrasse, *Notolabrus tetricus*. *Marine and Freshwater Research* **52**:311-322.
- Smith, D. C. 1994. Blue warehou. Page 360 in R. D. J. Tilzey, editor. *The South East Fishery: a scientific review with particular reference to quota management*. Bureau of Resource Sciences, Parkes.
- Smith, D. C., I. Montgomery, K. P. Sivakumaran, K. Krusic-Golub, K. Smith, and R. Hodge. 2003. The fisheries biology of bluethroat wrasse (*Notolabrus tetricus*) in Victorian waters. FRDC Report No. 97/128. Marine and Freshwater Institute, Victoria, Australia.
- Smith, H. K. 1983. Fishery and Biology of *Nototodarus gouldi* (McCoy, 1888) in western Bass Strait. *Memoirs of the National Museum Victoria* **44**:285-290.
- St Hill, J. L. 1996. Aspects of the biology of southern sea Garfish, *Hyporhamphus malanochir*, in Tasmanian waters. University of Tasmania, Hobart.
- Stark, K. E. 2008. Ecology of the arrow squid (*Nototodarus gouldi*) in southeastern Australian waters: a multi-scale investigation of spatial and temporal variability. University of Tasmania, Hobart.
- Steer, M. A., N. A. Moltschaniwskyj, and F. C. Gowland. 2002. Temporal variability in embryonic development and mortality in the southern calamary *Sepioteuthis australis*: a field assessment. *Marine Ecology Progress Series* **243**:143-150.
- Steer, M. A., N. A. Moltschaniwskyj, D. S. Nichols, and M. Miller. 2004. The role of temperature and maternal ration in embryo survival: using the dumpling squid *Euprymna tasmanica* as a model. *Journal of Experimental Marine Biology and Ecology* **307**:73-89.
- Stevens, J. D. and H. F. Hansfeld. 1982. Age determination and mortality estimates on an unexploited population of Jack mackerel *Trachurus declivis* (Jenyns, 1841) from south-east Australia. Report No 148., CSIRO Marine Laboratories, Cronulla.
- Stewart, J., J. Hughes, J. McAllister, J. M. Lyle, and M. MacDonald. 2011. Australian salmon (*Arripis trutta*): population structure, reproduction, diet and composition of commercial and recreational catches. Fisheries Final Report Series No. 129. Industry & Investment NSW.
- Sutton, C. P., D. J. MacGibbon, and D. W. Stevens. 2010. Age and growth of greenback flounder (*Rhombosolea tapirina*) from southern New Zealand. New Zealand Fisheries Assessment Report 2010/48. NIWA, Wellington.
- Thresher, R. E., C. H. Proctor, J. S. Gunn, and I. R. Harrowfield. 1994. An evaluation of electron-probe microanalysis of otoliths for stock identification of nursery areas in a southern temperate groundfish, *Nemadactylus macropterus* (Cheilodactylidae). *Fishery Bulletin* **92**:817-840.
- Tilzey, R. D. J. 1994. *The South East fishery: a scientific review with particular reference to quota management*. Bureau of Resources Sciences, Australia.
- Tilzey, R. D. J., M. Zann-Schuster, N. L. Klaer, and M. J. Williams. 1990. *The South East Trawl Fishery: biological synopses and catch distributions for seven major commercial fish species*.
- Tracey, S. R. and J. M. Lyle. 2005. Age validation, growth modeling and mortality estimates for striped trumpeter (*Latris lineata*) from south-eastern Australia: making the most of patchy data. *Fishery Bulletin* **103**:169-182.
- Tracey, S. R., J. M. Lyle, G. Ewing, P., K. Hartmann, and A. Mapleston. 2013. Offshore recreational fishing in Tasmania 2011/12. Institute for Marine and Antarctic Studies, Hobart.
- Tracey, S. R., J. M. Lyle, and M. Haddon. 2007a. Reproductive biology and per-recruit analyses of striped trumpeter (*Latris lineata*) from Tasmania, Australia: implications for management. *Fisheries Research* **84**:358-367.
- Tracey, S. R., A. J. Smolenski, and J. M. Lyle. 2007b. Genetic structuring of *Latris lineata* at localized and transoceanic scales. *Marine Biology* **152**:119-128.
- Triantafillos, L. 2004. Effects of genetic and environmental factors on growth of southern calamary, *Sepioteuthis australis*, from southern Australia and northern New Zealand. *Marine and Freshwater Research* **55**:439-446.

- Triantafillos, L. and M. Adams. 2001. Allozyme analysis reveals a complex population structure in the southern calamary *Sepioteuthis australis* from Australia and New Zealand. *Marine Ecology Progress Series* **212**:193-209.
- Triantafillos, L., G. D. Jackson, M. Adams, and B. McGrath Steer. 2004. An allozyme investigation of the stock structure of arrow squid *Nototodarus gouldi* (Cephalopoda:Ommastrephidae) from Australia. *ICES Journal of Marine Science* **61**:829-835.
- Uozumi, Y. 1998. Fishery biology of arrow squids, *Nototodarus gouldi* and *N. sloanii* in New Zealand Waters. *Bulletin of the National Research Institute of Far Seas Fisheries* **35**:1-111.
- van den Enden, T., R. W. G. White, and N. G. Elliott. 2000. Genetic variation in the greenback flounder *Rhombosolea tapirina* Günther (Teleostei, Pleuronectidae) and the implications for aquaculture. *Marine and Freshwater Research* **51**:23-33.
- Ward, T. M., O. Burnell, A. Ivey, J. Carroll, J. Keane, J. Lyle, and S. Sexton. 2015. Summer spawning patterns and preliminary Daily Egg Production Method survey of Jack Mackerel and Australian Sardine off the East Coast. South Australian Research and Development Institute (Aquatic Sciences).
- Wayte, S. E. 2013. Management implications of including a climate-induced recruitment shift in the stock assessment for jackass morwong (*Nemadactylus macropterus*) in south-eastern Australia. *Fisheries Research* **142**:47-55.
- Webb, B. F. 1976. Aspects of the biology of jack mackerel *Trachurus declivis* (Jenyns) from south east Australian waters. *Tasmanian Fisheries Research* **10**:1-17.
- Welsford, D. C. 2003. Early life-history, settlement dynamics and growth of the temperate wrasse, *Notolabrus fucicola* (Richardson 1840), on the east coast of Tasmania. PhD thesis. University of Tasmania, Hobart.
- Williams, H. and G. Pullen. 1986. A synopsis of biological data on the jack mackerel *Trachurus declivis* Jenyns. Tasmanian Department of Sea Fisheries Technical Report 10. 34 p.
- Wolf, B. 1998. Update on juvenile banded morwong in Tasmania. *Fishing Today* **11**:30.
- Woodhams, J., S. Vieira, and I. Stobutzki. 2012. Fishery status reports 2011. Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra.
- Woodhams, J., S. Vieira, and I. Stobutzki. 2013. Fishery status reports 2012. Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra.
- Ziegler, P. E. 2012. Fishing tactics and fleet structure of the small-scale coastal scalefish fishery in Tasmania, Australia. *Fisheries Research* **134-136**:52-63.
- Ziegler, P. E., M. Haddon, and J. M. Lyle. 2006. Sustainability of small-scale, data-poor commercial fisheries: Developing assessments, performance indicators and monitoring strategies for temperate reef species. Tasmanian Aquaculture and Fisheries Institute, Hobart.
- Ziegler, P. E., J. M. Lyle, and M. Haddon. 2008. Tasmanian Scalefish Fishery-2007. Hobart.
- Ziegler, P. E., J. M. Lyle, M. Haddon, and G. P. Ewing. 2007a. Rapid changes in life-history characteristics of a long-lived temperate reef fish. *Marine and Freshwater Research* **58**:1096-1107.
- Ziegler, P. E., J. M. Lyle, G. T. Pecl, N. A. Moltschanivskyj, and M. Haddon. 2007b. Tasmanian Scalefish Fishery-2006. Tasmanian Aquaculture and Fisheries Institute, Hobart.

Appendix 1- Common and scientific names of species from catch returns

Common name	Scientific name	Common name	Scientific name
Alfonsino	<i>Beryx</i> spp.	Sardine	Fam. Clupeidae
Anchovy	Fam. Engraulidae	Rays bream	Fam. Bramidae
Atlantic salmon	<i>Salmo salar</i>	Redbait	<i>Emmelichthys nitidus</i>
Australian Salmon	<i>Arripis</i> spp.	Red fish	Fam. Berycidae
Barracouta	<i>Thyrsites atun</i>	Red Mullet	<i>Upeneichthys</i> spp.
Boarfish	Fam. Pentacerotidae	Silverfish	Fam. Atherinidae
Bream	<i>Acanthopagrus butcheri</i>	Snapper	<i>Pagrus auratus</i>
Butterfish	Spp unknown	Stargazer	Fam. Uranoscopidae
Cardinal fish	Fam Apogonidae	Sweep	<i>Scorpiis</i> spp
Cod deep sea	<i>Mora moro</i>	Tailor	<i>Pomatomus saltatrix</i>
Cod, bearded rock	<i>Pseudophycis barbata</i>	Thetis fish	<i>Neosebastes thetidis</i>
Cod, red	<i>Pseudophycis bachus</i>	Trevalla, white	<i>Seriola caerulea</i>
Cod, unspec.	Fam. Moridae	Trevally, silver	<i>Pseudocaranx dentax</i>
Dory, john	<i>Zeus faber</i>	Trout, rainbow	<i>Oncorhynchus mykiss</i>
Dory, king	<i>Cyttus traversi</i>	Trumpeter, bastard	<i>Latridopsis forsteri</i>
Dory, mirror	<i>Zenopsis nebulosus</i>	Trumpeter, striped	<i>Latris lineata</i>
Dory, silver	<i>Cyttus australis</i>	Trumpeter, unspec.	Fam. Latridae
Dory, unspec.	Fam. Zeidae	Warehou, blue	<i>Seriola brama</i>
Eel	<i>Conger</i> spp.	Warehou, spotted	<i>Seriola punctata</i>
Flathead	Fam Plactycephalidae	Whiptail	Fam. Macrouridae
Flounder	Fam. Pleuronectidae	Whiting	Fam. Sillaginidae
Garfish	<i>Hyporhamphus melanochir</i>	Whiting, King George	<i>Sillaginoides punctata</i>
Gurnard	Fam. Triglidae & Fam. Scorpaenidae	Wrasse	<i>Notolabrus</i> spp.
Gurnard perch	<i>Neosebastes scorpaenoides</i>	'Commonwealth' spp	
Gurnard, red	<i>Chelidonichthys kumu</i>	Blue grenadier	<i>Macruronus novaezelandiae</i>
Hardyheads	Fam. Atherinidae	Gemfish	<i>Rexea solandri</i>
Herring cale	<i>Odax cyanomelas</i>	Hapuka	<i>Polyprion oxygeneios</i>
Kingfish, yellowtail	<i>Seriola lalandi</i>	Oreo	Fam. Oreosomatidae
Knifejaw	<i>Oplegnathus woodwardi</i>	Trevalla, blue eye	<i>Hyperoglyphe antartica</i>
Latchet	<i>Pterygotrigla polyommata</i>	Tunas	
Leatherjacket	Fam. Monacanthidae	Albacore	<i>Thunnus alalunga</i>
Ling	<i>Genypterus</i> spp.	Skipjack	<i>Katsuwonus pelamis</i>
Luderick	<i>Girella tricuspidata</i>	Southern bluefin	<i>Thunnus maccoyii</i>
Mackerel, blue	<i>Scomber australasicus</i>	Tuna, unspec.	Fam. Scombridae
Mackerel, jack	<i>Trachurus declivis</i>	Sharks	
Marblefish	<i>Aplodactylus arctidens</i>	Shark, angel	<i>Squatina australis</i>
Morwong, banded	<i>Cheilodactylus spectabilis</i>	Shark, blue whaler	<i>Prionace glauca</i>
Morwong, blue	<i>Nemadactylus valenciennesi</i>	Shark, bronze whaler	<i>Carcharhinus brachyurus</i>
Morwong, dusky	Fam. Cheilodactylidae	Elephantfish	<i>Callorhynchus milii</i>
Morwong, grey	<i>Nemadactylus douglasii</i>	Shark, gummy	<i>Mustelus antarcticus</i>
Morwong, jackass	<i>Nemadactylus macropterus</i>	Shark, saw	<i>Pristophorus</i> spp.
Morwong, red	Fam. Cheilodactylidae	Shark, school	<i>Galeorhinus galeus</i>
Morwong, unspec.	Fam. Cheilodactylidae	Shark, seven-gilled	<i>Notorynchus cepedianus</i>
Mullet	Fam. Mugilidae	Shark, spurdog	Fam. Squalidae
Nannygai	<i>Centroberyx affinis</i>	Cephalopods	
Perch, magpie	<i>Cheilodactylus nigripes</i>	Calamari	<i>Sepioteuthis australis</i>
Perch, ocean	<i>Helicolenus</i> spp.	Cuttlefish	<i>Sepia</i> spp.
Pike, long-finned	<i>Dinolestes lewini</i>	Octopus	<i>Octopus</i> spp.
Pike, short-finned	<i>Sphyaena novaehollandiae</i>	Squid, Gould's	<i>Nototodarus gouldi</i>

Appendix 2- Data restriction and adjustments

There have been a number of administrative changes that have affected the collection of catch and effort data from the fishery. The following restrictions and adjustments have been applied when analysing the data as an attempt to ensure comparability between years, especially when examining trends over time.

Tasmanian logbook data

i) Correction of old logbook landed catch weights

Prior to 1995, catch returns were reported as monthly summaries of landings. With the introduction of a revised logbook in 1995, catch and effort was recorded on a daily basis for each method used. Since catch data reported in the old general fishing return represent landed catch, it has been assumed to represent processed weights. For example, where a fish is gilled and gutted, the reported landed weight will be the gilled and gutted and not whole weight. By contrast, in the revised logbook all catches are reported in terms of weight and product form (whole, gilled and gutted, trunk, fillet, bait or live). If a catch of a species is reported as gilled and gutted then the equivalent whole weight can be estimated by applying a standard conversion factor³.

Without correcting for product form, old logbook and revised logbook catch weights are not strictly compatible. In an attempt to correct for this and provide a 'best estimate', a correction factor was calculated using catch data from the revised logbook and applied to catches reported in the old logbook. A species based ratio of the sum of estimated whole weights (adjusted for product form) to the sum of reported catch weights was used as the correction factor (Lennon 1998).

ii) Effort Problems

Records where effort (based on gear units, Table 2.1) was zero or null, or appeared to be recorded incorrectly (implausible), were flagged. The catch was included in catch summaries but the records were not included in gear unit effort and catch rate calculations. These records were, however, used in calculating days fished and daily catches.

iii) Vessel restrictions

In all analyses of catch and effort, catches from six vessels (four Victorian based and two Tasmanian based) have been excluded. These vessels were known to have fished consistently in Commonwealth waters and their catches of species such as Blue Warehou and ling tended to significantly distort catch trends. In fact, all four Victorian vessels and one of the Tasmanian vessels ceased reporting on the Catch and effort logbooks in 1994. With the introduction of the South East Fishery Non-Trawl logbook (GN01) in 1997, the remaining Tasmanian vessel ceased reporting fishing activity in the Tasmanian logbook.

³ Conversion factors to whole weights are 1.00 for whole, live or bait; 2.50 for fillet; 1.50 for trunk; and 1.18 for gilled and gutted.

Commonwealth logbook data:

Commonwealth logbook data from Australian Fisheries Management Authority was included in the analyses so that the assessment reflected all catches from Tasmanian waters.

(i) Area restrictions

Commonwealth logbook records were only included if the catch was taken in fishing blocks adjacent to Tasmania and the maximum depth of the fishing operation was less than 200 m. These conditions were applied to all records except where striped or Bastard Trumpeter were caught. All records that included catches of these species were included for analysis, because these species are managed under Tasmanian jurisdiction in all waters adjacent to Tasmania.

Fishing blocks adjacent to land and used in the analyses (refer Fig. A1) include:

3C2, 3D1, 3F1, 3F2, 3G1, 3G2, 3C4, 3D3, 3F4, 3G3, 3G4, 3H3, 3H4, 4C2, 4D1, 4D2, 4E1, 4G2, 4H1, 4H2, 4D4, 4E3, 4E4, 4F4, 4G3, 4G4, 4H3, 4H4, 5D2, 5E2, 5F1, 5F2, 5H1, 5D4, 5E3, 5H3, 6E1, 6H1, 6E3, 6G4, 6H3, 7E1, 7E2, 7G1, 7G2, 7H1, 7E4, 7F3, 7F4, 7G3.




Fig. A1 Numbers for fishing blocks used in calculation of catch figures.

(ii) Duplicate records

A number of records in Commonwealth logbooks had matching records (fisher, date, gear type) in the Tasmanian database. Such records were examined individually and decisions made as to whether it was more appropriate to keep the Tasmanian record, the Commonwealth record or both. In most situations the Tasmanian logbook entry was kept and the Commonwealth record excluded. The only exceptions were records with extra information in the Commonwealth record, e.g. catch of a Commonwealth species that was not recorded in the Tasmanian logbook.

Appendix 3- Annual Tasmanian scalefish production

Table A.1. Annual Tasmanian scalefish and cephalopod production (whole weight in tonnes) by financial year since 1990/91 based on Catch and effort logbooks and Commonwealth (GN01, GN01A and SSJF) logbook returns. Note that the data includes Commonwealth catches of Striped Trumpeter, Bastard Trumpeter and Gould's Squid in State waters. Data of the most recent years may be incomplete due to late reporting.

Species	90/91	91/92	92/93	93/94	94/95	95/96	96/97	97/98	98/99	99/00	00/01	01/02	02/03	03/04	04/05	05/06	06/07	07/08	08/09	09/10	10/11	11/12	12/13	13/14
Scalefish (excl. small pelagics)																								
Australian Salmon	815.9	651.9	867.0	878.8	682.1	413.2	287.3	475.7	384.7	363.7	485.0	462.1	407.2	167.2	336.5	254.2	114.9	171.5	338.8	372.3	203.5	189.4	331.3	37.7
Barracouta	351.5	268.3	205.4	59.6	25.2	19.3	53.8	65.2	27.6	25.0	15.1	132.1	65.5	85.2	97.3	60.1	25.0	10.7	11.7	7.6	5.0	4.0	1.1	1.1
Boarfish	7.2	9.4	7.6	10.1	9.1	7.3	10.0	6.2	3.2	2.5	3.6	5.5	3.6	4.3	3.6	5.0	4.9	4.6	2.4	2.7	1.9	3.4	2.1	1.0
Cod	10.0	11.3	11.6	14.5	12.7	18.6	12.8	9.4	9.6	8.7	3.7	3.0	2.2	2.1	1.6	2.0	2.6	2.0	3.3	2.6	2.8	2.3	2.0	2.0
Flathead	165.3	118.1	98.8	121.4	91.1	57.9	51.8	62.7	50.5	60.2	63.1	52.1	40.8	31.2	74.7	91.9	59.3	73.1	52.6	77.3	55.7	62.1	40.2	31.1
Flounder	44.0	36.8	31.8	27.3	27.1	33.4	29.4	29.7	25.2	18.6	12.3	13.0	10.9	14.9	14.7	10.9	12.5	6.3	4.9	5.2	5.2	4.0	2.0	2.1
Garfish	80.9	80.1	82.3	82.9	69.3	56.2	91.6	83.0	101.7	91.2	81.4	87.8	92.5	66.2	85.5	89.3	45.8	23.6	60.7	49.3	43.2	53.0	51.5	34.3
Gurnard	20.5	19.0	19.3	19.3	14.0	13.5	10.4	9.1	7.0	9.5	7.4	5.3	9.7	6.8	6.1	5.1	5.6	4.4	2.6	1.5	2.1	1.1	1.1	0.6
Leatherjacket	12.2	14.0	13.1	23.3	27.7	14.5	12.6	13.3	12.9	16.5	16.7	16.6	13.7	14.8	10.4	8.5	8.7	4.7	5.2	3.0	2.8	2.2	2.4	2.9
Ling	5.1	13.6	30.0	41.6	33.2	15.0	13.3	8.3	4.3	1.8	1.2	0.9	0.4	0.8	0.7	0.4	0.4	0.4	0.1	0.1	0.1	0.1	0.1	1.2
Marblefish	0.2	0.9	0.3	1.0	1.8	3.5	5.6	3.0	2.6	4.2	4.0	4.4	3.1	0.6	1.1	0.5	2.0	2.2	1.0	0.5	0.2	0.2	0.3	0.2
Morwong, banded	7.0	6.9	39.2	145.5	105.8	86.7	79.0	72.6	42.4	33.8	39.2	53.7	56.0	46.4	45.6	54.3	50.3	52.6	36.9	44.6	40.9	40.3	37.9	34.2
Morwong, jackass	136.9	111.9	83.2	117.6	63.1	27.1	18.7	33.2	17.5	15.7	13.1	14.8	14.4	16.3	17.5	13.1	11.6	4.3	5.2	5.9	3.2	2.8	1.5	1.0
Morwong, other	3.8	5.6	5.2	13.9	8.1	5.4	7.4	7.4	6.3	1.4	0.6	1.4	1.9	1.2	1.8	1.3	1.3	2.2	1.4	1.2	0.9	0.7	0.7	0.6
Mullet	31.2	22.2	26.2	19.5	23.8	10.8	11.2	16.0	14.5	21.0	13.7	12.1	7.3	7.5	5.1	7.5	4.5	2.4	1.5	1.9	2.2	1.7	9.2	2.9
Other	140.2	110.4	97.4	102.0	62.0	26.3	24.2	24.6	17.1	13.6	10.4	10.6	30.1	21.3	25.0	26.8	13.8	8.7	31.6	4.7	2.5	3.5	2.3	2.1
Pike	10.5	9.5	11.1	12.7	18.8	14.0	18.3	21.6	12.6	14.0	13.3	20.4	17.9	18.2	8.9	13.9	16.3	15.4	12.5	11.2	11.7	10.8	8.2	9.9
Trevally	20.6	13.6	12.0	8.3	21.6	8.4	6.0	5.4	6.5	2.7	1.6	4.7	5.9	3.4	3.7	6.3	3.5	8.6	4.5	3.8	1.9	2.1	5.4	4.1
Trumpeter, bastard	63.3	37.2	34.0	54.8	50.8	60.1	51.8	40.4	47.5	36.2	25.7	23.9	21.0	23.2	18.5	23.4	19.9	18.1	16.0	10.5	9.8	9.4	9.4	8.2
Trumpeter, striped	74.5	58.2	52.7	56.5	72.4	58.3	79.4	78.1	99.0	95.0	45.5	39.9	36.5	36.9	23.9	19.0	16.9	9.9	10.6	10.8	19.7	20.9	13.0	10.3
Trumpeter, unspec.	0.7	0.0	0.0	0.4	0.1	0.0	0.1	0.6	3.3	0.0				0.0				0.1	0.0	0.1	0.1	0.0	0.1	0.1
Warehou, blue	257.6	317.6	187.7	250.1	205.4	82.3	128.4	187.6	272.2	186.1	34.2	66.4	49.3	27.5	19.1	20.0	28.4	24.6	26.6	37.5	10.7	3.8	8.5	5.2
Warehou, other	0.7	0.4	4.2	8.8	3.4	14.6	15.6	4.2	1.0		0.0	0.1	0.2	0.1	0.8	0.1	0.0	0.1	0.6	0.2	0.0	0.0	0.2	0.0
Whiting	124.2	152.3	84.3	97.9	81.4	25.5	39.6	48.3	30.6	31.7	42.7	40.1	39.9	55.5	38.3	28.3	40.2	39.6	35.0	27.0	34.6	16.1	16.5	37.2
Wrasse	57.2	71.7	97.3	142.4	178.0	83.4	110.1	100.0	90.7	85.4	88.4	92.3	72.0	75.1	100.1	92.9	109.3	71.4	66.3	72.0	71.7	66.0	63.7	62.4
Total scalefish (excl. small pelagics)	2441	2140	2101	2310	1888	1155	1168	1405	1290	1138	1021	1163	1002	726.8	940.6	834.8	597.5	561.2	731.9	753.8	532.3	499.8	610.7	292.5

Table A.1 Continued. Whole weight in tonnes by financial year

Species	90/91	91/92	92/93	93/94	94/95	95/96	96/97	97/98	98/99	99/00	00/01	01/02	02/03	03/04	04/05	05/06	06/07	07/08	08/09	09/10	10/11	11/12	12/13	13/14
Small pelagic species																								
Mackerel, jack	6.1	11.1	32.8	48.4	39.7	26.2	19.3	19.7	59.8	13.3	8.6	19.4	19.4	41.1	12.8	6.8	2.6	8.0	788.7	917.0	35.7	56.4	0.2	0.4
Mackerel, other (blue)	3.0	2.1	0.3	8.5	5.7	2.0	1.3	1.0	0.5	2.1	0.1	0.0	0.1		0.5	0.5	0.2	5.3	0.2	0.3	0.8	0.1	1.9	4.2
Sardine/Anchovy	0.1	0.0	3.8	14.6	12.1	6.6	4.3	15.4	2.8	1.7	3.2	0.7		0.3	0.8			10.2	13.2	0.4				0.1
Redbait	0.0	0.7	0.0	0.8	0.1	0.1			4.0					3.4	1.0	1.4	0.3	156.4	441.4	121.6	15.0	0.1	0.1	
Total small pelagic species	9.2	13.9	36.9	72.3	57.6	34.9	24.8	36.1	67.0	17.1	11.9	20.1	19.5	44.8	15.2	8.7	3.1	179.9	1243	1039.3	51.6	56.7	2.1	4.6
Cephalopod																								
Calamari, southern	8.2	7.5	5.8	9.7	12.6	33.0	19.0	26.6	94.4	84.6	76.6	104.8	108.8	86.8	114.2	44.5	80.2	80.0	77.7	51.1	54.9	50.8	63.9	67.5
Cuttlefish	0.5	0.7	0.0	1.1	0.8	0.2	0.3	0.2	0.0	0.0	0.0	0.7	2.4	1.0	0.2	0.4	0.1	0.3	0.3	0.1	0.1	0.1	0.2	0.1
Octopus pallidus ⁴	21.2	21.6	31.3	29.8	30.5	58.7	30.2	34.1	61.2	43.2	50.5	31.6	54.5	56.6	70.4	89.9	90.2	92.6	64.5	65.1	92.3	57.9	124.8	100.0
Squid, Gould's	35.1	7.2	7.0	7.7	8.6	5.7	7.8	12.9	79.7	480.5	39.7	2.4	1.9	2.1	2.6	1.8	5.8	1.4	16.8	121.3	131.2	516.6	1071.8	0.0
Total cephalopod	65.0	37.0	44.1	48.3	52.5	97.7	57.3	73.8	235.3	608.4	166.8	139.4	167.7	146.5	187.4	136.6	176.3	174.2	159.3	237.5	278.5	625.3	1260.6	167.6
Sharks⁵																								
Elephantfish						58.0	48.9	21.4	14.7	17.0	16.7	18.4	16.5	10.2	7.6	5.7	9.0	1.8	1.5	2.4	1.3	2.6	1.9	1.4
Gummy shark						750.5	543.8	348.6	113.4	109.6	53.9	23.5	14.2	24.7	41.6	12.4	12.6	12.6	9.4	9.8	9.3	7.5	7.9	5.9
Draughtboard shark						0.0	0.0	0.0	0.0	0.0	0.0	1.7	0.7	1.0	0.8	1.3	1.2	0.4	0.3	0.2	0.3	0.0	0.1	0.0
Sawshark						127.4	74.4	29.2	6.8	3.4	12.3	21.4	20.4	20.6	23.5	5.9	3.4	0.3	0.1	0.1	0.1	0.4	0.0	0.0
School shark						252.1	171.5	71.7	31.5	11.3	1.7	2.2	1.4	7.0	2.6	0.6	1.8	0.9	0.7	1.8	1.4	1.9	2.1	1.0
Seven-gilled shark						6.1	4.9	6.1	1.9	10.3	16.3	18.8	7.4	11.5	8.4	3.8	3.7	0.5	2.3	1.1	1.4	1.1	0.8	0.7
Other shark						26.4	16.1	11.3	6.8	6.5	4.8	5.8	3.6	3.2	1.1	0.6	2.2	0.9	0.7	0.3	0.8	0.5	0.5	0.7
Total sharks						1220	859.6	488.2	175.1	158.2	105.7	91.9	64.3	78.3	85.6	30.4	34.0	17.4	15.1	15.7	14.7	14.1	13.3	9.8

⁴ Octopus catches in NWC and NEC were assumed to be *Octopus pallidus* (excluding bycatch in rock lobster traps). Catches can also include *Octopus tetricus*.

⁵ Since 2001/02, shark catches have been reported in Commonwealth logbooks. Tasmania has jurisdiction of all shark species inside 3nm except gummy and school shark, and fishers are on bycatch possession limits for all species. Includes bycatch from the rock lobster fishery (separate reporting since 2007/08).

Appendix 4- Annual catch, effort and number of vessels by fishing methods

Table A.2 Total annual catch, effort and number of vessels by fishing methods.

Effort units are defined in Table 1.1. * Catch data not shown where five or fewer vessels involved.

Gear	Year	Catch(t)	Effort#	Days fished	Vessels
Graball net	95/96	348.0	223553	5437	257
	96/97	383.3	231140	5186	232
	97/98	446.3	231412	5249	216
	98/99	493.3	166505	4689	204
	99/00	359.7	152144	4169	203
	00/01	173.4	86838	3187	186
	01/02	196.0	71109	3303	180
	02/03	231.0	85628	3395	168
	03/04	189.8	69189	2904	160
	04/05	154.4	53965	2491	137
	05/06	170.2	51591	2402	123
	06/07	170.0	56487	2543	132
	07/08	161.3	59308	2425	115
	08/09	127.1	47468	1977	95
09/10	128.7	47049	2022	94	
10/11	98.1	45324	1770	79	
11/12	100.6	40150	1583	71	
12/13	102.9	40390	1538	60	
13/14	99.20	37811	1366	56	
Small mesh net	95/96	38.7	10971	285	19
	96/97	27.0	7965	260	14
	97/98	21.8	7875	246	17
	98/99	31.2	7772	282	14
	99/00	22.7	6232	210	15
	00/01	20.8	8170	256	14
	01/02	24.7	9863	259	11
	02/03	22.9	10297	284	11
	03/04	23.0	7254	228	11
	04/05	15.3	5982	220	13
	05/06	21.7	5890	191	11
	06/07	16.4	7144	202	11
	07/08	15.3	6447	183	10
	08/09	9.5	4817	152	7
09/10	9.4	4089	134	7	
10/11	7.1	2281	98	7	
11/12	10.9	4127	152	8	
12/13	17.4	4784	189	10	
13/14	10.0	3270	146	10	
Dip-net	95/96	5.6	320	83	5
	96/97	24.2	1518	364	10
	97/98	37.9	1903	449	21
	98/99	43.6	2784	579	29
	99/00	29.4	2319	505	35
	00/01	22.8	1430	371	27
	01/02	24.8	1561	387	27
	02/03	18.7	1259	337	20
	03/04	25.6	1557	374	19
	04/05	27.4	1521	305	16
	05/06	39.1	2167	376	18
	06/07	22.6	1308	244	18
	07/08	15.2	1057	227	17
	08/09	16.4	1229	287	14
09/10	18.0	1376	270	14	
10/11	16.0	958	227	10	
11/12	22.5	1187	220	9	
12/13	23.2	1205	172	7	
13/14	16.9	1115	147	7	

Table A.2 (continued)

Gear	Year	Catch(t)	Effort#	Days fished	Vessels
Beach seine	95/96	469.2	1086	559	53
	96/97	351.7	1355	685	50
	97/98	520.9	1206	582	44
	98/99	441.7	872	398	40
	99/00	422.9	901	430	33
	00/01	528.4	789	373	31
	01/02	572.2	1070	495	30
	02/03	490.7	1063	511	35
	03/04	238.1	1282	458	31
	04/05	397.0	975	368	27
	05/06	308.4	653	304	25
	06/07	140.6	528	234	25
	07/08	266.3	630	263	17
	08/09	382.6	620	282	12
	09/10	410.3	473	251	15
10/11	225.3	359	200	13	
11/12	203.7	380	184	11	
12/13	334.3	321	162	11	
13/14	45.02	251	148	12	
Danish seine	95/96	68.2	474	163	2
	96/97	70.7	360	116	1
	97/98	93.1	456	133	1
	98/99	67.6	375	94	1
	99/00	74.5	515	139	2
	00/01	101.4	589	152	2
	01/02	77.7	491	145	2
	02/03	68.1	354	129	3
	03/04	74.1	278	127	2
	04/05	95.9	282	108	2
	05/06	104.9	418	132	3
	06/07	85.8	475	157	3
	07/08	103.2	482	162	3
	08/09	72.9	387	134	3
	09/10	97.0	557	181	4
10/11	81.1	385	155	5	
11/12	68	260	94	3	
12/13	47	327	112	5	
13/14	59.2	369	116	3	
Purse seine	95/96	35.2	418	185	11
	96/97	30.4	336	153	10
	97/98	41.8	319	154	7
	98/99	76.9	246	150	8
	99/00	33.7	244	123	10
	00/01	24.6	224	104	4
	01/02	21.3	216	91	5
	02/03	18.7	139	76	4
	03/04	14.8	68	45	3
	04/05	17.6	130	70	5
	05/06	16.0	122	60	4
	06/07	8.1	86	41	4
	07/08	527.8	117	121	5
	08/09	1486.3	195	153	3
	09/10	1033.0	170	99	4
10/11	53.9	32	32	3	
11/12	70.5	63	60	3	
12/13	33.4	94	90	3	
13/14	7.13	23	23	2	

Table A.2 (continued)

Gear	Year	Catch(t)	Effort#	Days fished	Vessels
Hand line	95/96	74.3	16964	1612	147
	96/97	94.3	21542	1893	135
	97/98	97.5	21076	1702	145
	98/99	88.2	17668	1278	127
	99/00	87.8	16688	1439	134
	00/01	74.2	13585	1541	130
	01/02	87.3	15527	1603	138
	02/03	72.2	15025	1552	125
	03/04	76.4	15610	1411	127
	04/05	100.5	19953	1803	123
	05/06	82.6	20247	1884	116
	06/07	107.3	22745	2139	128
	07/08	92.0	19985	2032	119
	08/09	80.5	17930	1741	95
	09/10	84.1	17259	1601	90
10/11	90.1	18006	1647	89	
11/12	80.4	15925	1634	82	
12/13	75.6	12348	1377	76	
13/14	74.8	13069	1244	66	
Drop-line	95/96	19.9	438	158	31
	96/97	30.0	433	203	27
	97/98	24.7	539	222	42
	98/99	31.8	666	309	38
	99/00	30.8	385	291	48
	00/01	15.8	382	248	36
	01/02	12.8	220	258	35
	02/03	18.8	264	350	43
	03/04	19.4	378	281	51
	04/05	14.1	351	219	31
	05/06	9.3	185	204	33
	06/07	7.1	259	137	28
	07/08	3.0	39	55	19
	08/09	3.1	45	41	18
	09/10	2.4	35	25	7
10/11	4.7	58	45	14	
11/12	6.5	98	59	10	
12/13	8.4	156	76	11	
13/14	4.2	112	45	6	
Fish trap	95/96	41.8	8264	1401	66
	96/97	57.2	10710	1796	66
	97/98	49.9	9870	1875	71
	98/99	53.7	10657	1559	56
	99/00	56.1	11030	1637	62
	00/01	54.3	9356	1548	68
	01/02	49.0	6098	1278	62
	02/03	38.2	6177	1246	58
	03/04	48.0	6308	1414	58
	04/05	46.7	7409	1222	54
	05/06	44.6	12302	1421	54
	06/07	44.2	11001	1328	47
	07/08	27.9	9761	916	44
	08/09	13.8	5316	568	29
	09/10	9.5	2257	341	23
10/11	9.0	2696	408	22	
11/12	8.6	2934	338	18	
12/13	6.9	1585	188	15	
13/14	8.7	1649	266	13	

Table A.2 (continued)

Gear	Year	Catch(t)	Effort#	Days fished	Vessels
Squid-jig	95/96	10.2	5389	125	23
	96/97	5.7	640	77	14
	97/98	15.2	4381	211	18
	98/99	89.8	10200	613	53
	99/00	150.3	39240	989	64
	00/01	66.5	13173	793	53
	01/02	85.2	12544	925	65
	02/03	91.8	19220	1228	68
	03/04	69.8	15764	1223	73
	04/05	104.8	22362	1424	79
	05/06	35.4	11223	767	59
	06/07	74.4	14105	1204	67
	07/08	77.7	16220	1406	52
	08/09	71.0	15339	1079	35
	09/10	47.5	14865	920	37
10/11	49.3	12932	866	27	
11/12	42.1	11378	753	23	
12/13	65.7	11380	826	26	
13/14	52.2	11154	790	22	
Spear	95/96	14.1	1403	368	21
	96/97	19.3	1853	464	27
	97/98	16.8	1981	483	40
	98/99	19.8	1812	452	38
	99/00	19.3	2233	475	25
	00/01	14.4	1586	355	22
	01/02	13.1	1296	279	19
	02/03	10.3	1366	247	22
	03/04	10.5	1446	289	22
	04/05	13.5	1609	357	24
	05/06	7.9	1009	271	22
	06/07	15.4	1414	362	20
	07/08	9.8	957	255	21
	08/09	5.6	886	181	14
	09/10	5.0	663	168	14
10/11	4.0	626	178	15	
11/12	5.0	622	167	12	
12/13	3.3	489	126	12	
13/14	3.7	339	105	7	
Troll	95/96	19.6	3497	358	55
	96/97	62.1	9755	600	88
	97/98	76.2	13318	685	83
	98/99	46.1	9307	480	73
	99/00	39.7	6184	420	51
	00/01	36.5	7913	440	48
	01/02	141.1	18664	1033	93
	02/03	82.8	8510	676	75
	03/04	87.7	8995	793	89
	04/05	87.7	6797	758	77
	05/06	63.4	5019	491	54
	06/07	30.3	4664	341	53
	07/08	13.3	2648	172	30
	08/09	20.0	3007	175	28
	09/10	12.3	2487	137	22
10/11	10.9	2358	110	19	
11/12	10.4	1970	93	11	
12/13	5	1622	72	10	
13/14	5.4	2078	79	12	

Table A.2 (continued)

Gear	Year	Catch(t)	Effort#	Days fished	Vessels
Automatic squid-jig	95/96	0.4	3000	2	1
	96/97	1.3	14560	14	3
	97/98	2.5	3040	8	1
	98/99	49.7	133728	66	3
	99/00	386.9	899711	369	13
	00/01	28.1	280200	115	10
	01/02	0.9	10220	7	3
	02/03	0.6	9000	5	1
	03/04	0.1	3000	2	1
	04/05	0.1	500	1	1
	05/06	-	-	-	-
	06/07	681.9	395762	155	10
	07/08	44.4	55657	21	3
	08/09	42.4	22966	12	3
	09/10	118.5	3329	68	5
10/11	2.3	98	7	5	
11/12	159.5	2454	60	7	
12/13	974.5	10640	214	7	
13/14	0	0	0	0	
Hand collection	95/96	7.0	1198	433	6
	96/97	2.5	298	154	5
	97/98	4.2	261	214	4
	98/99	11.5	918	267	4
	99/00	7.2	947	228	3
	00/01	5.7	1008	272	3
	01/02	19.5	1185	318	4
	02/03	6.6	625	160	3
	03/04	5.9	678	184	3
	04/05	3.1	400	128	3
	05/06	2.0	226	97	4
	06/07	4.5	439	201	5
	07/08	8.4	571	207	10
	08/09	1.0	122	52	4
	09/10	0.8	124	55	4
10/11	4.9	378	120	8	
11/12	1.6	177	99	3	
12/13	2.1	228	115	3	
13/14	4.0	352	134	6	